

Objectives & Significance

Unravelling the evolutionary history of the Milky Way has been a long-standing problem in contemporary astrophysics, and understanding this history will have significant ramifications for our insight into how other galaxies form and evolve. The critical step will be to make precise measurements of the fundamental properties of stars – the building blocks and the source of galactic chemical evolution – something we currently have achieved only in the solar neighbourhood. Traditional near-field cosmology, or galactic archaeology, studies have been limited by the indirect way the crucial stellar parameters, age and mass, can be inferred for distant stars.

The Kepler and CoRoT missions have revealed an exciting new opportunity for progress: red giants are high-amplitude oscillators, and the analysis of their light curves can yield asteroseismically-determined radii, masses and ages. This discovery has tremendous potential for expanding our view into how the Galaxy formed and evolved. However, the data from these pioneering studies is insufficient: they sample only a limited portion of galactic real estate, and a lack of well-described selection criteria for the stellar sample limits our ability to faithfully compare theoretical models with observations (Chaplin et al. 2011).

K2 provides a fantastic opportunity to overcome this hurdle. *The thrust of this proposal is to combine K2 data from a carefully-selected stellar sample with ambitious ground-based observations, and include state-of-the-art stellar and galactic modelling calculations.* The project will complement traditional spectroscopic and photometric diagnostics with those derived from the rapidly emerging field of asteroseismology. Applying asteroseismology to obtain the radii, masses and ages of a large ensemble of stars across the Galaxy will enable us to determine much more accurately the fundamental physical processes that led to the present-day Milky Way.

Asteroseismology is based on the elegant notion that the oscillations of an object can teach us about its physical properties; the power of this approach in the case of the Sun has been strongly demonstrated (Christensen-Dalsgaard, 2002). Convective surface motion excites sound waves, causing the star to oscillate in many modes simultaneously (Fig. 1). The frequencies depend on the physical properties of the stellar interior, such as density, temperature and composition.




Fig.1: Model of oscillation mode in a star.

Measurements of stellar oscillation frequencies combined with spectroscopy, which provides stellar surface properties, therefore yield crucial information about mass, radius, and age that have proven very difficult to obtain for field stars in the past.

With the launch of the space missions CoRoT and particularly Kepler, red giant asteroseismology went wholesale. Crucial for the project, the detection of oscillations in large ensembles of intrinsically luminous and hence distant red giant stars (Hekker et al. 2011, Stello et al. 2013) enable the probing of stellar populations in the Galaxy (Miglio et al. 2009, 2013).

‘Galactic archaeology’ is a field in astrophysics that studies galaxy evolution through the fossil evidence from the stellar relics of ancient star formation (Freeman & Bland-Hawthorn, 2002). The field has recently gained significant traction thanks to ambitious programs aimed at measuring the chemical composition of more than a million stars, including the Australian GALAH survey led by Freeman and Bland-Hawthorn and the US-led APOGEE survey.

Central to galactic archaeology – and the overall aim of this proposal, is to address the following critical questions facing our current paradigms for galactic evolution: (1) is the halo made entirely of debris of small accreted galaxies, or does it include stars born in-situ in the Milky Way, (2) how is the thick disk formed, and (3) what does the age–metallicity and velocity–metallicity relations look like at large distances from the Sun. A recent gathering of world experts in galaxy evolution and asteroseismology (Sesto, Italy, 2013) has singled out the need to obtain precise stellar ages – which asteroseismology can provide – as the most important issue in the field of galactic archaeology. *Resolution of the above issues is likely to have profound consequences for a wide range of fields in astrophysics.*

Approach & Methodology

We base our approach on observing a large number of colour-magnitude selected red giants to probe the Galaxy far beyond the solar neighbourhood. This proposal is part of the continuing K2 Galactic Archaeology Program, which was initiated at Campaign-0 and allocated 8629 stars during Campaign-1, and 5138 stars in Campaign-2 – the largest number of allocated targets to a single proposal. It is our intention to make similar proposals for future K2 fields in order to probe galactic directions not probed before taking advantage of K2’s unique ‘360-degree view’.

Target selection & number of targets

The goal of galactic archaeology is to understand both the initial conditions (where and when each star is born) and the complex processes governing the evolution (the interaction of each star with the rest of the Galaxy) leading to the present day Milky Way. This problem can be formulated by four stellar observables: position, velocity, age, and metallicity – the latter a tracer of galactic radius at birth. In the ideal case one would want to sample the entire parameter space defined by our four observables in order to translate that into a complete picture of the origin of the stars and their galactic evolution. This would require of order 1 million stars (Freeman & Bland-Hawthorn 2002). If the number of stars available is much less, as in our case with K2, one needs to invoke additional assumptions such as exploring a limited set of parameters or using parametric forms for certain physical relationships.

With the K2 Galactic Archaeology Program we aim at projecting the full parameter space into few dimensions such that we can explore key aspects of galactic evolution, in particular the age-metallicity distribution and age-velocity distribution and their variation in galactic radius, R , and height above the plane, z . For this we require about 5,000 (oscillating red giant) stars each at 10 unique R and z locations within the galaxy, which breaks down to about 5,000 red giants per K2 campaign. These numbers are based on our expected observational uncertainties on age (20% or 0.08 dex) and metallicity (0.1dex), and the need to sample the parameter range spanned by the Galaxy, age: 1–13.7 Gyr (14 bins), metallicity: $-2 < [\text{Fe}/\text{H}] < 0.5$ (25 bins) by at least 15 stars per bin, which translates to a S/N ratio above 3.8 (Poisson statistics). Each campaign (1–9) field has different giant yields ranging 50–90%; hence 5,500 to 10,000 targets need to be observed per campaign to reach the required 5,000 red giants.

Our targets are selected based purely on their 2MASS colour and magnitude. We select stars having $J-K > 0.5$, $V > 9$ and sort them by V magnitude (brightest at top) derived as $V = K + 2*((J-K) + 0.14) + 0.382*\exp(2(J-K-0.2))$, approximately ranging $9.5 < Kp < 14.5$. Stars that are already observed spectroscopically by APOGEE, RAVE, or Gaia-ESO (typically 100-500

per campaign) are placed at the top of the list. Our colour-magnitude cuts ensure a well-described selection function, aid efficient ground-based follow up, and constitute the sweet spot for where we expect to detect and resolve oscillations in red giants without pushing the K2 pixel budget by highly saturated stars. We refer to the summary tables and corresponding figures in the Target List section for quantitative details.

Towards the faint end, our selection will serendipitously include a significant and increasing number of dwarfs of great value to all surveys looking for planets around cool stars, as well as stellar rotation, activity, and binarity. Hence, although we advocate for a minimum of order 5-10 thousand targets per campaign, “going down” our target list even further, boosts targets suited for other science goals while adding relatively few targets unique to our project. This ensures a reproducible selection function for all investigations using these data – a win-win situation.

Generating light curves

We will generate light curves using an aperture photometry pipeline developed for the K2 mission. The software, written in Python, performs dynamic automated aperture mask selection, background estimation and subtraction, and positional decorrelation to remove as best as possible effects due to spacecraft micro-slews and pointing jitter. Development and testing of the pipeline is ongoing based on K2 engineering data and should be fully functional by the time C4 and C5 data are available. Since spacecraft pointing performance are likely to vary slightly based on the particular field of view, we will optimize the pipeline for each campaign. The light curves will be (post-) processed to further reduce effects such as outliers, jumps or thermal drifts, following the automated methods described in Garcia et al. (2011) and Mathur et al. (in prep).

Measuring asteroseismic parameters and surface properties

The measurements of the large frequency separation, $\Delta\nu$, and frequency of max power, ν_{\max} , will be done using the SYD (Huber et al. 2009) and A2Z (Mathur et al. 2010) time series analysis pipelines. We further anticipate participation from other groups within KASC WG8 using independent methods. From these analyses, we create a consolidated list of results, which will be fed into the next stage of the analysis – the grid modelling. The full potential of the grid modelling will be reached when combined with data on stellar surface temperature and metallicity. Through our team’s strong links to large ground-based surveys, these additional data are already available to us for a significant fraction of our targets, and will for the remainder be added as top priority on the target lists of the APOGEE, GALAH and SAGA surveys.

Grid modelling

To obtain estimates of the stellar radius, mass and age we will use the RADIUS pipeline (Stello et al. 2009). RADIUS uses seismic scaling relations and large grids of stellar models to estimate fundamental stellar parameters based on $\Delta\nu$, ν_{\max} , T_{eff} and $[\text{Fe}/\text{H}]$. Through collaboration with KASC WG8, we will engage a larger set of grid modelling pipelines, which will help us obtain robust results and inform typical systematic errors. As demonstrated in Stello et al. (2009) (see also Huber et al. 2012, Silva Aguirre et al. 2012 and Chaplin et al. 2013), stellar radii can be obtained to the 2–3% level and mass to 7% from grid modelling. From preliminary work we expect to obtain ages to about 15–20%, but we aim to improve on that with complementary theoretical work and improved calibrations of asteroseismic scaling relations. These efforts will build on our current analysis of the sensitivity of red giant models to changes in the input physics, and on our continued calibration of seismic scaling relations using open clusters,

eclipsing binaries, and interferometry (Stello et al. 2011, Brogaard et al. 2012, Huber et al. 2012, and White et al. 2013). The PI and collaborators are heavily involved in a comprehensive analysis of the sensitivity of red giant models to changes in the input physics. They are also involved in the ongoing APOKASC project combining spectroscopic and asteroseismic data in the original Kepler field.

Modelling the Galaxy

We will use GALAXIA (Shama et al. 2011) and TRILEGAL (Girardi et al. 2005), two state-of-art galaxy synthesis tools, to create theoretical populations that represent the targeted stars. Comparing observations with theory will allow us to test the currently assumed physics, especially the age–metallicity and age–velocity relations that underpin the galactic models. The comparison will use our seismic results on radius, mass, and age, and T_{eff} and $[\text{Fe}/\text{H}]$ drawn mainly from APOGEE and GALAH, and where applicable, from other surveys such as RAVE (Steinmetz et al. 2006), Gaia-ESO (Gilmore et al. 2012), LAMOST (Cui et al. 2012), and SAGA (Casagrande et al. 2014). We will include kinematics and stellar parameters from ESA’s Gaia mission when they become available.

Our asteroseismic analyses will determine the distances to tens of thousands of stars, which will be a major asset. Initially, we will do this using our seismically inferred stellar radii; combined with T_{eff} , this will provide a luminosity estimate at the 10% level, and distances to a precision of ~5%. After the end of the grant we expect to improve our age estimates using complementary distances from Gaia. This will be particularly important for investigating the build-up of the thin and thick disks and the halo at the very early evolution of the Galaxy. With our results, we aim to distinguish between the oldest accreted stars and stars formed in-situ in the inner halo.

Impact & Relevance

The project will dramatically advance our knowledge about the Milky Way. With major ramifications for the evolution theory of all spiral galaxies, the research will provide a self-consistent picture of the stellar age distribution for tens of thousands of stars in the Milky Way, revealing the first precise timeline for how the Galaxy formed and evolved.

Understanding how galaxies evolve helps to inform our perception of the cosmos, from a cultural to a psychological level. The work outlined in this proposal addresses NASA’s Strategic Goal 1, as defined in the 2014 Strategic Plan: “Expand the frontiers of knowledge, capability, and opportunity”, including Objective 1.6: “Discover how the universe works, explore how it began and evolved, and search for life on planets around other stars.” In addition, the proposed project directly addresses Section 3.2 of NASA’s 2013 Astrophysics Roadmap: “Archaeology of the Milky Way and Its Neighbors.”

The highly-sampled and high-precision photometric data that K2 can provide are crucial for the project. No other current mission can provide data that will allow us to detect oscillations in large numbers of red giants. Our program takes full advantage of K2’s unique ‘360 degree’ field-of-view capability by probing the Galaxy along the ecliptic over the course of subsequent K2 campaigns; this will provide data from vastly different stellar populations within all four main components of the Milky Way – the thin and thick disks, the halo and the bulge. Our strong links to large industrial scale spectroscopy surveys makes our targets top priority for intense ground-

based follow up. These aspects ensure that the resulting data set arising from this program will be of great use for the broader community in future studies, leading to a long-lasting legacy of the K2 mission.

Plan of work

The work outlined in this proposal is a large multi-national coordinated effort. It comprises the primary research for PI Stello, who recently won a 4-year Fellowship sponsored by the Australian Research Council to work specifically on the asteroseismology of red giants to perform galactic archaeology investigations.

Key milestones

- We expect the first phase of the project (generating light curves) to complete four months from the date pixel data are made public [D+4mth]. At the time of pixel data release we anticipate our pipeline is ready and tested on previous K2 data, and light curves should be generated with minimal campaign-specific modifications required.
- Subsequent completion of time series analysis to extract $\Delta\nu$, and ν_{\max} , of all red giants in our sample including compilation of the consolidated values [D+6mth].
- K2 targeting is an identified priority for APOGEE and GALAH, and we expect to be able to have spectra for approved targets in fields 4 and 5 by the time the K2 data are obtained.
- Completion of the following grid modelling, including consolidating the results from the different grids [D+8mth].
- Compare estimated radii, masses and ages with population synthesis models of the Galaxy (initial results expected after four months) [D+12mth].

Management structure

The collaboration behind this proposal comprises large parts of KASC Working Group 8 (red giants), asteroSTEP (a collaboration aimed at merging expertise on galaxy and stellar evolution with asteroseismic techniques), APOKASC (a collaboration between KASC and APOGEE focussed on merging asteroseismic data from the old Kepler field with ground-based infrared spectroscopy), and GALAH (a large project on the AAT in Australia to obtain spectra of up to 1mio stars in the southern hemisphere). In addition to the PI and Co-Is, the 28 participants that have committed time to this project are listed in the proposal's Program Specific Data point 7.

Dennis Stello (PI): is the overall coordinator of the project. Stello was among the early pioneers in applying seismology to study red giants. He was co-creator of the SYD pipeline designed to extract seismic parameters from Kepler data, and he led the first demonstration of asteroseismic grid modelling techniques. He will use his existing software to determine the seismic inferred stellar properties. As part of the KASC WG8 chairing group, Stello will liaise with KASC on efforts where expertise and capabilities can be efficiently utilised within the broader WG8 community including independent time series analyses and grid modelling, which has been common practise during similar projects aimed at the Kepler and CoRoT fields.

Marc Pinsonneault (Co-I): is the primary link to the APOGEE consortium, which will provide surface temperatures and metallicities for the stars in our northern fields. These data will be

crucial for the grid modelling process to obtain precise estimates of the stellar properties. Marc Pinsonneault led similar work within APOKASC (Pinsonneault et al. 2014).

Ken Freeman (Co-I): is the senior leader of the GALAH project on the AAT. GALAH serves a similarly important role as APOGEE, but with an emphasis on stars towards the south. In collaboration with his team, he will be responsible for providing stellar surface temperatures and metallicities to the project.

Andrea Miglio (Co-I): is the leader of the asteroSTEP collaboration, and has been a pioneer in using seismic measurements to probe stellar populations in the Galaxy. He will be responsible for coordinating comparisons between our observed stellar properties and those found from simulations of the Galaxy in close collaboration with Co-I Sanjib Sharma and Leo Girardi.

Derek Buzasi (Co-I): is the brainchild of the resurrection of NASA's WIRE satellite and developed the software to generate light curves from its star tracker. He will lead the work on creating the K2 light curves of our targets. This will be in coordination with KASC-wide efforts.

Savita Mathur (Co-I): is the co-developer of the A2Z time series analysis code built to measure Δv , and v_{\max} of stars observed by Kepler and CoRoT data, and she has been a central part of the team that has generated corrected light curves for KASC in the past. She will work closely with Rafael Garcia (Saclay) and Co-I Buzasi to provide light curves optimised for seismic purposes, and she will take part in the time series analysis.

Sanjib Sharma (Co-I): is the developer of the GALAXIA population synthesis generator. He will work with Co-I Ken Freeman, and Joss Bland-Hawthorn (University of Sydney) to optimize target selection for the GALAH spectroscopic survey and with Co-I Miglio on comparing our stellar data with galaxy models.

Data sharing plan

We will make the light curves of our targets available. This will potentially constitute a large fraction of all stars observed by K2, with direct benefit for other science goals including the search for planets around cool dwarf stars, investigations of binary stars, and the study of stellar rotation. Creating the light curves will be coordinated with KASC, and as for past KASC data products, they will be available on KASOC and offered to become publicly available at MAST.


In addition, we will provide a catalogue of seismic quantities (Δv , v_{\max}), spectroscopic surface properties (T_{eff} and $[\text{Fe}/\text{H}]$) and estimated radii and masses of the red giants. APOGEE is part of the Sloan Digital Sky Survey, and APOGEE spectroscopic data will be either provided as part of a value-added catalogue with the K2 data or as part of the 2016 Sloan public data release. Similarly, the GALAH data used in this project will either be provided as part of our catalogue or made public beforehand through GALAH specific programs. The asteroseismic data from the K2 fields, which complement current observations from the Kepler Cygnus field and CoRoT, will form a legacy data set for galactic archaeology. Combined with the forthcoming data from Gaia, the results from these efforts will be a true goldmine, allowing exploration of core areas of galactic evolution theory including the chemical and dynamical evolution of the Milky Way.

Target list

The target list is not provided in this document due to the large number of targets proposed. Instead we give a more informative target list summary table for each campaign and show the corresponding plots of the table columns.


Campaign-4

V~Kp+0.35	N_target	N_giant	Giant_fraction
< 10	157	149	0.95
< 11	525	498	0.95
< 12	1435	1305	0.91
< 13	3451	2829	0.82
< 14	7914	5223	0.66
< 15	17410	8008	0.46



Campaign-5

V~Kp+0.35	N_target	N_giant	Giant_fraction
< 10	211	206	0.98
< 11	595	577	0.97
< 12	1478	1389	0.94
< 13	2881	2506	0.87
< 14	5267	3844	0.73
< 15	9937	5167	0.52



References

- Brogaard et al. 2012, A&A, 543, 106
Casagrande et al. 2014, ApJ, 787, 110
Chaplin et al. 2011, Science, 332, 213
Chaplin et al. 2013, ApJS, 210, 1
Christensen-Dalsgaard, 2002, RvMP, 74, 1073
Cui et al. 2012, RAA, 12, 1197
Freeman & Bland-Hawthorn, 2002, ARA&A, 40, 487
Garcia et al. 2011, MNRAS, 414, 6
Gilmore et al. 2012, The Messenger, 147, 25
Girardi et al. 2005, A&A, 436, 895
Hekker et al. 2011, MNRAS, 414, 2594
Huber et al. 2009, CoAst, 160,74
Huber et al. 2012, ApJ, 760, 32
Mathur et al. 2010, A&A, 511, 46
Miglio et al. 2009, A&A, 503, 21
Miglio et al. 2013, MNRAS, 429, 423
Pinsonneault et al. 2014, ApJ (in press)
Shama et al. 2011, ApJ, 730, 3
Silva Aguirre et al. 2012, ApJ, 757, 99
Steinmetz et al. 2006, AJ, 132, 1645
Stello et al. 2009, ApJ, 700, 1598
Stello et al. 2011, ApJ, 737, 10
Stello et al. 2013, ApJ, 765, 41
White et al. 2013, ApJ, 751, 36