


**National Committee for Space Science Decadal Plan
Final report of the Planetary Science Working Group**

26 March 2007



Members of the NCSS-DP Planetary Sciences Working Group:

Dr. Marc Norman, chair (Australian National University)
Prof. Malcolm Walter (Macquarie University)
Prof. Brad Carter (University of Southern Queensland)
Prof. Ross Taylor (Australian National University)
Dr. David Nelson (Curtin University of Technology)
Dr. Vickie Bennett (Australian National University)
Dr. Graziella Caprarelli (University of Technology Sydney)
Dr. Jon Clarke (Geoscience Australia)
David Stegman (Monash University)
Dr. Sergey Vladimirov (University of Sydney)

With submissions from:

Dr. Jeremy Bailey (Macquarie University)
Dr. Trevor Ireland (Australian National University)
Prof. Robert Pidgeon (Curtin University)

Executive Summary

Planetary science addresses fundamental problems such as the origin of the solar system and whether life exists on other planets. This type of research has the potential to generate transformative science that can change the way we view the world and our selves.

An increasing number of nations are planning spacecraft missions to the planets, their satellites and smaller bodies of the Solar System. In the next ten years the USA and China will be well advanced in their plans to return humans to the Moon, and Europe, India and Japan will have a strong presence in space. Spacecraft missions will visit Mars, Venus, Mercury, the Moon, comets, asteroids, and the outer planets. Earth-based satellites will contribute to an improved appreciation of the space environment and how it impacts life on Earth. The advent of new players in Asia and Europe opens opportunities for international cooperation and Australian participation in these high-profile activities.

Planetary science offers a cost effective means for Australia to participate in this prestigious international arena. Space engages the public's imagination and can inspire young people in their choice of careers. Students gain a strong foundation in mathematics and fundamental sciences, and the use of advanced technology, thereby fostering creativity and promoting a culture of innovation.

Despite the complete absence of a space program, Australian planetary scientists have developed a strong international reputation in planetary geology, cosmochemistry, and planetary geophysics. Many individual Australian scientists are at the forefront of current research into the origin and evolution of the Moon and the early Earth, meteorite impacts, astrobiology, and the evolution of planetary interiors. Australian scientists are, however, disadvantaged by a lack of coordination at the national level.

Science Goals:

This draft report of the Planetary Sciences Working Group outlines topics and questions that represent high priority research areas for the Australian planetary science community in the coming decade. Much of the research described here is focused on the origin and early evolution of the terrestrial planets and astrobiology. Because many Australian planetary scientists are employed in related disciplines (e.g. earth science, astronomy, physics) several of the research priorities identified here have synergies with strategic research goals articulated by the National Committees for Astronomy and Earth Sciences.

The Working Group identified the following planetary science themes expected to generate major conceptual advances in the coming decade:

1. How did the planets of our Solar System form?
2. How and why do planetary surfaces and atmospheres change over geological time?
3. What is the role of meteorite impacts in planetary evolution?
4. Is there life elsewhere in the solar system and beyond?

Recommendations:

The Planetary Science Working Group recommends that the NCSS support the establishment of a virtual centre for planetary science. The goal of such a centre would be to provide more effective links between existing concentrations of planetary scientists within Australia.

The Working Group also makes a strong recommendation that the NCSS support the establishment of an organization charged with long term planning, co-ordination and implementation of Australian space science and technology programmes at the national level. Mechanisms to more fully integrate Australian space scientists with spacecraft missions through technology development and advanced data processing should be developed.

Table of Contents

1. Introduction

- 1a. What is Planetary Science?
 - 1b. National Benefits
 - 1c. Challenges and Recommendations
 - 1d. Education and Public Outreach
-

2. Key Science Questions

Theme #1: How and over what timescales did planets form from the solar nebula?

- 2a. Evolution of the Solar Nebula: Precursors to planets
 - 2a-1. Nucleosynthesis and stellar contributions to the Solar Nebula
 - 2a-2. Dynamics of Dusty Plasmas
 - 2a-3. The composition of the Sun
 - 2b. Accretion and Early Differentiation of Terrestrial Planets
-

Theme #2: How and why do planetary surfaces, atmospheres, and climates change over time?

- 2c. Planetary geodynamics
 - 2c-1. Surface environments
 - 2c-2. Planetary atmospheres
 - 2c-3. Australian analogues for planetary surfaces
-

Theme #3: What is the role of meteorite and asteroid impacts in planetary evolution?

- 2d. The Impact of impacts
-

Theme #4: Is there life elsewhere in the solar system or beyond?

- 2e. Habitable Planets and the Uniqueness of Earth
 - 2e-1. Origin of Life
 - 2e-2. Search for Earth-like planets
-

3. Submissions from Australian planetary science research groups

- 3a. Geological Society of Australia Specialist Group in Planetary Science.
- 3b. Planetary Astronomy Group, Australian Centre for Astrobiology

1. Introduction

1a. What is Planetary Science?

Planetary science will be one of the most exciting and rapidly expanding frontier research areas of the coming decade. The recent discovery of more than 200 extrasolar planets and the continuing exploration of our Solar System through spacecraft missions and studies of extra-terrestrial materials provides powerful stimuli for understanding the geological and astrophysical processes that create planetary systems and shape the surfaces and interiors of planetary bodies.

Planetary science is inter-disciplinary and international in scope, linking earth scientists, astronomers, physicists, chemists, biologists, engineers, and mathematicians working independently on topics as diverse as the origin of our Solar System, global climate change, and development of new technologies for laboratory analysis and remote sensing. We bring samples of other planets and extinct stars into our laboratories, and we map the surfaces and probe the interiors of other planets using data obtained by spacecraft. Our goal is to understand the origin of the Solar System and our place in it.

The knowledge base among planetary scientists in Australia is vast but somewhat disjointed and less effective than it might be, in part because of lack of coordination at the national level. National support will drive breakthrough science, grow new technologies, and enhance Australia's international prestige.

The Planetary Sciences Working Group identified four major themes that comprise some of the most fertile areas for research in the coming decade. These are:

1. How and over what timescales did the planets of the Solar System form?
2. How do planetary surfaces, atmospheres, and climates change over time?
3. What is the role of meteorite and asteroid impacts in planetary evolution?
4. Is there life elsewhere in the solar system or beyond?

These questions provide a framework within which achievable, high-impact planetary science outcomes over the coming decade have been identified. These key science questions and supporting material are presented in the following section of this report.

1b. National benefits

Planetary science in Australia is directed primarily toward **National Research Priority 3, Breakthrough Science and Promoting an Innovation Culture and Economy**. A solid foundation of knowledge-based skills is a critical component of a national comparative advantage. National benefits accrue through an increase in Australia's stock of knowledge, a primary driver in the information economy, and the production of graduates skilled in science and technology

National benefits also accrue through the production of graduates skilled in mathematics, fundamental science, and technology. Both of these outcomes are recognised as vital to the country's success in using, developing, and adopting new technologies, as well as making informed decisions on the increasing number of science-based issues that confront modern society.

Planetary science research showcases **Australian scientific and technical leadership** in a field of global interest, thereby enhancing the international visibility and national prestige. Space science enjoys widespread public interest and support, and research such as this has the potential to inspire the imagination of young people considering careers in science and technology.

The coming decade will see a flotilla of spacecraft missions to Mars, Venus, Mercury the Moon, asteroids, comets and the outer planets by European, Chinese, Indian, and Japanese space agencies as well as NASA. This broadly based international effort has generated renewed enthusiasm for planetary science worldwide. Australia will benefit from participation in these **high-profile international endeavours** where possible. Research support for planetary science offers one of the most direct and cost effective means by which Australia can contribute to this prestigious international arena.

1c. Challenges and Recommendations

The Working Group identified a number of significant challenges and structural impediments to effective and efficient involvement of Australian planetary scientists with national and international collaborators. These include (1) the dispersed nature of Australian planetary scientists across the country, (2) the lack of a national focus for planetary science research and mission development, (3) structural restrictions on access to mission data and new technology, (4) emphasis on short-term funding cycles that effectively preclude significant involvement in mission planning. These constraints place artificial limits on the abilities of Australian scientists to function at the highest international level and reduce the potential national benefits that planetary science can deliver.

The Planetary Science Working Group makes three **Recommendations**:

1. We recommend that the NCSS Decadal Plan support the establishment of a virtual centre in planetary science. Australian planetary scientists are widely distributed across the country and there is currently no central focus for the broad range of planetary science being conducted in Australia. One of the goals of such a virtual centre would be to provide more effective opportunities for collaboration between existing concentrations of planetary scientists in Australia. Such a centre will help build a **critical mass** of Australian planetary scientists, and improve collaborations between Australian planetary scientists, earth scientists, and astronomers. An essential goal is to improve links between science and technology groups within Australia to identify and enhance mission concepts and instrumentation, and leverage interactions with industry and defence.

2. We recommend that the Decadal Plan identify research areas that would benefit from acquisition and development of new instruments, experimental facilities, and data processing capabilities linked to research components of the plan.

3. We recommend the development of appropriate organizations and institutional vehicles for the long term planning, co-ordination and implementation of space science and technology development programmes at the **national level** to improve the delivery of professional and educational opportunities, and benefits to society.

1d. Outreach and Public Education

Planetary science is accessible, rigorous, and timely. In many ways it is an ideal vehicle for public education and outreach because it:

- * Engages people's imaginations and their natural curiosity about the world in which they live.
- * Motivates an aspirational society to compete and achieve at the highest international level.
- * Connects people with their home planet and their place in it.
- * Provides a natural basis for education integrating physical sciences, mathematics, and engineering,
- * Bridges scientific disciplines.
- * Increases the global knowledge base at the macro and micro level.
- * Encourages discourse and provides common grounds for action among scientists, educators, industry partners, and policy makers.

2. Key Science Questions

In this section we describe some of the areas of planetary science in which major progress will be made in the coming decade and in which Australia is well placed to make significant contributions. This section is organized according to the four themes identified in the Introduction, followed by a description of Key Science Questions and supporting materials.

Theme #1: How and over what timescales did planets of the Solar System form?

2a. Evolution of the Solar Nebula: Precursors to planets

Apart from light chemical elements (such as H, He and Li) created during the Big Bang by spallation reactions, the chemical elements of which our solar system is composed were produced principally within the interiors of stars. We know that a variety of exploding stars contributed newly synthesised ejecta to the interstellar molecular cloud from which our solar system was ultimately created. The dust and gas that condensed from the stellar ejecta formed the starting materials for our solar system. This section focuses on the origin of these starting materials and the processes that transformed the dust and gas of an interstellar cloud into the planets and central star that makes up our Solar System.

2a-1. Nucleosynthesis and stellar contributions to the Solar Nebula

Key Science Question #2a-1:

* What was site of formation, initial solar system abundance, and spatial distribution of short-lived radioactive isotopes such as ^{26}Al , ^{60}Fe , and ^{10}Be in the solar nebula?

Significance of Key Science Question #2a-1:

These data are necessary to advance our understanding of the astrophysical environment of the inner solar system and timescales of planetary accretion relative to formation other nebular objects to the next level. The initial abundance of ^{60}Fe in the solar system would pinpoint either a supernova or AGB-type star as the source of short-lived radionuclides. This in turn would constrain the relative abundances of other short-lived nuclides from this source, allowing other contributions such as Solar Energetic Particle interactions to be quantified

Background

The chemical composition of our solar system was determined by the mixing of nuclides synthesized in different proportions within many different stellar sources over almost 10

billion years. Geochemical studies of primitive meteorites have identified radioactive decay products from about a dozen short-lived (with half-lives ≤ 100 million years) radionuclides that were present at the early stages of nebular condensation. Studies of these short-lived radioactive decay systems in meteorites, terrestrial and lunar samples have enabled establishment of a remarkably detailed high-resolution chronology of some of the key early condensation, accretion and planetary differentiation events in our solar system. Short-lived radionuclides have also been detected in relatively recent (3 million year old) deep-sea sediments by accelerator mass spectrometry. Such radioactive fallout from space may have played a significant role in Earth's mass extinction and climate history.

The origins of these nuclides and their importance as heat sources during planetary accretion and differentiation is a topic of vigorous research. There are three alternative explanations for the existence of short-lived radionuclides in early solar system materials: (i) they were derived from an exploding novae, supernovae or a red giant star in the vicinity and within 2 Ma of formation of our solar system (ii) they were derived by stellar nucleosynthesis reactions within many different stars in the galactic neighborhood at the time of solar system formation (i.e. Background Uniform Production), (iii) some radionuclides may have been synthesized by spallation interactions with energetic particles from the early and unstable (T Tauri) sun. The presence of radionuclides with very short half-lives (e.g. ^{41}Ca , which has a half-life of $\sim 10^5$ yrs) requires that a stellar nucleosynthesis event occurred within 2 Ma of the onset of formation of our solar system. The shock waves that originated from this exploding star could have triggered the rapid collapse of a more slowly evolving molecular cloud of interstellar dust and gas nearby, thus initiating the formation of our own Sun and solar system. The nature of the last stellar source that contaminated our solar system's precursor molecular cloud with freshly synthesized elements may be inferred from the relative abundance of short-lived nuclides in meteorites. Leading contenders are a thermally pulsing Asymptotic Giant Branch star, a Type Ia or Type II supernova, or a Wolf-Rayet star (a massive, short-lived star that ends its life as a supernova). An additional "galactic uniform production" contribution from earlier nucleosynthesis events may be required to account for abundances of the longer-lived radionuclides, and some of the short-lived nuclides can apparently be produced efficiently by spallation reactions with energetic solar particles.

Nucleosynthesis yields within stellar sites are currently not well constrained, principally due to uncertainties associated with key nuclear reaction rates. Fortunately, yields of several short-lived nuclides and the galactic uniform production rate may be determined by direct observation. Gamma rays are an important by-product of radioactive decay. Recent gamma-ray observations using NASA's RHESSI and ESA's INTEGRAL satellites have facilitated refinement of current quantitative models of stellar nucleosynthesis yields of some short-lived nuclides. Currently only a subset of the meteorite data have been adequately explained. A comprehensive model is needed that identifies the most plausible sources for short-lived nuclides in the solar nebula.

In addition to the isotopic record of extinct radionuclides preserved within meteorites, micron-sized grains that condensed from the gas phase in cooling outflows of stars such

as red giants and supernova explosions prior to formation of our solar system have also been identified in primitive meteorites. These pre-solar dust grains formed in the interstellar medium and have made their way into the cloud of gas and dust that eventually formed our Solar System. Although dust-sized particles would have been vaporized as the T-Tauri Sun formed and heated up, a small proportion of these grains survived intact, probably protected inside asteroids that have been sampled as meteorites.

Studies of meteorites for extinct nuclides and pre-solar grain, linked to satellite gamma-ray spectrometry on the interstellar medium, can be used to constrain the physical environment in which the solar nebula evolved, the extent of radial and vertical mixing within the nebula, the contributions of different stellar sources to the nebula, and the chronology of nebular evolution and planet formation chronology.

Experiments and facilities needed to address Key Science Question #2a-1:

Analytical methods have been developed using the Curtin SHRIMP ion microprobe and are currently being applied to determine the abundance ratios of short-lived nuclides that are diagnostic of nucleosynthesis processes within AGB stars and supernovae. This will enable identification of the stellar source of the short-lived nuclides detected within early solar system materials.

2a-2. Dynamics of Dusty Plasmas

Key Science Question #2a-2:

* What role did dusty plasma environments play in the early solar nebula?

Significance of Key Science Question #2a-2:

Dusty plasma environments in the early solar nebula may provide a unifying perspective on diverse phenomena such as the origins of poorly understood objects from the solar nebula, the processes that create planets from dust and gas, and the behaviour of the Sun. Critical factors for understanding the evolution of the solar nebula and formation of the planets such as the distribution of dust and gas, the origin of high-temperature transient events, and processes of planetary accretion are poorly understood. Significant new discoveries in these areas are expected over the coming decade.

Background

Dust is a common constituent in many space and astrophysical environments, including molecular clouds, proto-planetary nebulae, stellar outflows, and supernovae explosions. All of these environments contributed in one way or another to the formation of our Solar System, so understanding the formation and behaviour of dust in these environments is fundamental to understanding the origin of the Solar System.

The physical and chemical processes that act on cosmic dust and that lead to conversion of dust to planetary bodies will be one of the leading research topics in the coming decade. It has the potential to be one of the most synergistic areas of space science, linking cosmochemistry, meteoritics, and fundamental physics.

Of specific interest is the fact that dusty space environments often exist in the presence of plasma. Plasmas are conductive assemblies of charged particles, neutrals and fields that exhibit collective effects. Plasmas are the most common form of matter, comprising more than 99% of the visible universe. They permeate the solar system, interstellar and intergalactic environments. Plasma temperatures and densities range from relatively cool and tenuous (like aurora) to very hot and dense (like the central core of a star). Of primary significance for understanding the behaviour of dust in space environments is the fact that plasmas carry electrical currents and generate magnetic fields.

Low-temperature plasmas in space often contain massive and heavily charged dust grains. Plasmas containing small solid particles ranging in size from nanometers to micrometers are called 'dusty plasmas'. Dusty plasmas are common in space, occurring in such diverse environments as interstellar clouds, interplanetary dust, comets, planetary rings, and the Earth's magnetosphere. In addition, the study of the influence of dust in the Earth's ionosphere and atmosphere is an important area for space weather and environmental research. Because of charge redistribution, the presence of dust in a plasma can also strongly affect its collective and transport properties. It is therefore of interest to investigate the behavior of inhomogeneous dusty plasmas and dusty plasma environments in the early Solar system.

The presence of electrically charged dust in the early solar nebula may have strongly affected almost all aspects of nebular evolution and planetary accretion. For example, dust dynamics and transport, dust clouds and streams, may have controlled the initial stages of agglomeration of nanometer-size dust grains into larger planetesimals, a process that is poorly understood at present. The presence of charged dust also affects plasma collective processes such as wave propagation, plasma instabilities, which must have contributed to mixing and phase separation in the early nebula. The related physical phenomena have relevance across a wide range of space environments, from cometary comae and tails to planetary atmospheres and the asteroid belt, from circumsolar dust rings to the noctilucent clouds in the arctic troposphere. Thus investigations of the physics and chemistry of dusty plasmas may provide keys that unlock current mysteries surrounding such phenomena as the formation of high-temperature nebular objects (e.g., chondrules and refractory inclusions), accretion of planetesimals, and the distribution of dust, gas, and associated chemical species within the nebula.

Certain constituents of primitive meteorites record transient high-temperature environments in the solar nebula. For example, chondrules are round grains of silicate minerals that formed as molten droplets free-floating in space. Chondrules formed by rapid heating (within minutes or less) of solid precursor material to temperatures between 1500°C and 1900°C followed by a cooling within one to several hours. The astrophysical

environment and the energy source for chondrule formation are unknown but much current research is directed toward consideration of shock waves in the solar nebula as a possible mechanism for chondrule formation. Another class of high-temperature objects from the early nebula are the calcium-aluminum inclusions (CAIs). These are highly refractory objects composed of oxides and silicates of calcium, aluminum, and titanium that would be among the first phases to condense from a cooling plasma. The mass-dependent fractionation of Mg and Si isotopes in some CAIs are consistent with evaporation or condensation of silicates. In addition, however, many CAI's have peculiar fractionated isotopic compositions of refractory elements such as Ti, Sr, Ba, and Eu that cannot be accounted for by volatility. Alternatively, magnetic separation in stellar outflow environments in plasmas surrounding the young Sun might account for this non-mass dependent isotope fractionation.

These puzzling aspects of chondrules and CAIs may be explicable in terms of dusty plasma environments in the solar nebula. For example, the presence of charged dust grains causes asymmetries in the plasma that change the behavior of shock waves and the reconnection process. Dust causes long-wavelength oscillations in the magnetic field upstream or downstream of a shock and causes the current filaments to twist and rotate about each other like a giant catherine wheel, or to merge asymmetrically. The extent to which these types of phenomena can explain the observed characteristics of nebular objects such as chondrules and CAIs will require integrated theoretical and experimental studies of the physics and chemistry of plasma environments linked to observations of nebular objects found in primitive meteorites.

Experiments and facilities needed to address Key Science Question #2a-2:

Significant progress in this area will require new collaborations between the Australian space scientists investigating the physics of dusty plasma environments and the chemistry of early nebular objects. One possible mechanism might be the establishment of experimental facilities suitable for investigating the physics and chemistry of dusty plasma systems of specific relevance to the early solar system. For example, the structural and chemical compositions of dust particles created experimentally in high-temperature systems could be investigated using electron and ion microprobe facilities at the University of Sydney, Curtin University and the Australian National University, and using the Australian synchrotron facilities that will be available in 2007. The experimental facility might be located at the University of Sydney to leverage existing expertise in plasma research. New nano-sample handling facilities similar to those used for inter-planetary dust particles (i.e., ultramicrotome etc.) will be needed to conduct these experimental and analytical studies. In addition, new measurements of the ages and isotopic compositions of CAIs and new theoretical models for their formation will be needed to establish possible nebular environments responsible for mass-dependent and non-mass dependent isotopic fractionations.

2a-3. The composition of the Sun

Key Science Question #2a-3:

* What is the oxygen and carbon isotopic composition of the Solar System?

Significance of Key Science Question #2a-3:

One of the most puzzling aspects of cosmochemistry is the astonishingly wide range of oxygen isotopic compositions observed within primitive meteorites. This may reflect a systematic difference in oxygen isotopic composition of the planets relative to their host star. A significant difference in O-isotopic composition between the Sun and the planets between might arise if there is a difference in isotopic composition between the dust and gas of the primordial molecular cloud as might be generated by photochemical reactions. In this scenario, the planets would have been sourced entirely from the dust component, whereas the Sun obtains a substantial fraction of its oxygen from carbon monoxide gas. Alternatively, early nebular condensates may have formed in a unique astrophysical environment. Distinguishing between these possibilities and placing better constraints on the processes and physical environment of the solar nebula would significantly improve our understanding the formation of the Solar System.

Background

Oxygen is one of the most abundant elements in the solar system, yet components found in meteorites display a 5% range in $^{18}\text{O}/^{16}\text{O}$ and $^{17}\text{O}/^{16}\text{O}$. Much of this variability seems to be related to mixing, implying the existence of significant heterogeneity in the early Solar System. Two popular models to explain this variability and predict the bulk composition of the Solar System have been proposed. Because increasingly large solid bodies (e.g., asteroids, Moon, Mars, Earth) have similar ^{16}O abundances, the bulk solar system may have an oxygen isotopic composition similar to that of the terrestrial planets. Alternatively, early refractory inclusions that have been interpreted as solar condensates have elevated ^{16}O abundance, raising the possibility that the Sun, and therefore the Solar System, has a distinctive ^{16}O -rich composition. The first alternative would require formation of the refractory inclusions in a unique astrophysical environment, whereas the second possibility would require significant processing and heterogeneity in the nebula to create such strong compositional differences between the Sun and the planets. A better understanding of either possibility would significantly improve our understanding of nebular evolution.

Experiments and facilities needed to address Key Science Question #2a-3:

A direct measurement of solar compositions is available from materials that have been exposed to the solar wind. Atoms are implanted in to the top 100 nm of mineral grain

surfaces. Analytical methods for in-situ measurements of small amounts of implanted oxygen have been developed at the Australian National University, and have been applied to studies of lunar metal grain exposed to the solar wind in the lunar regolith. The oxygen isotopic composition measured in the lunar grains is enriched in ^{17}O and ^{18}O by $5.3 \pm 0.3\%$ relative to terrestrial oxygen. This is in good agreement with a new measurement of the solar photosphere that indicates $\delta^{18}\text{O} = +4 \pm 6\%$. The relatively large errors are consistent with a normal (terrestrial) oxygen isotopic composition, but do not support a ^{16}O -rich composition for the Sun. The GENESIS spacecraft mission to the Sun returned small amounts of solar wind implanted on ceramic and metal plates. Despite the hard landing of the spacecraft and subsequent terrestrial contamination of the collector plates, it may still be possible to extract meaningful information using SHRIMP ion microprobe technology. Additional measurements on lunar metal grains and GENESIS collector plates are needed to assess the significance of these results.

2b. Accretion and Early Differentiation of Terrestrial Planets

Key Science Question #2b-1:

* What were the timescales and processes that created internally structured planets from the dust and gas of the solar nebula?

Significance of Key Science Question #2b-1:

Recent results suggest that large-scale planetary processes such as melting and core formation were occurring on asteroids at the same time as the high-temperature nebular events recorded by chondritic meteorites. This is difficult to rationalise within the context of our current understanding of nebular evolution. Geochemical and petrologic studies of igneous rocks from the Moon, Mars, Earth, and differentiated asteroids will advance our understanding of the early dynamics of terrestrial planets, the structure of planetary interiors, and the origins of differentiated bodies in the Solar System. This will provide a direct comparison with timescales for disk evolution estimated by astronomical methods.

Background

The processes that built planets from the dust and gas solar nebula are poorly understood. Astronomers and planetary scientists now agree that instabilities in the solar nebula lead to rapid growth of km-size bodies km-size planetesimals, probably within 10,000-100,000 years after initial collapse. Some of these planetesimals remained cool; chondritic meteorites are samples of these primitive bodies. On other planetesimals, heat liberated by accretion and decay of short-lived radionuclides such as ^{26}Al and ^{60}Fe caused extensive melting and allowed cores of molten iron to separate from their silicate mantles. Basaltic lavas erupted to form a primary crust. The mechanical and thermal effects of collisions among early accreting planetesimals may have contributed to melting, phase separation, and redistribution of volatile elements within the planetesimals

Accretion of rocky planetesimals apparently was synchronous with some of the earliest datable events in the solar nebula as recorded by primitive meteorites. Large-scale melting and internal differentiation of proto-planetary bodies occurred on timescales of 10's of millions of years. This contrasts dramatically with the prevailing view of just a few decades ago in which planets heated slowly over billions of years by accumulation of internal heat from long-lived radioactive decay. Rapid timescales of planet formation implies dynamic accretion probably involving large collisions between protoplanetary bodies and internal heating by short-lived radioactive heat sources.

Ages determined from terrestrial and lunar rocks and meteorites using a diverse array of radioactive decay schemes combined with theoretical models can now provide constraints on the timing of major planetary events including accretion, core formation and crustal growth on the Earth, Moon, Mars, and igneous asteroids. Spurred by analytical advances and the acquisition of larger planetary sample suites, an increasingly detailed view of the timing and processes responsible for early planetary differentiation is emerging.

Experiments and facilities needed to address Key Science Question #2b-1:

Laboratory studies of planetary materials are needed to establish a high-resolution absolute timescale for the early geological evolution of the inner solar system and the processes responsible for early planetary differentiation. These studies would entail geochemical analysis of radiogenic isotopes, trace element abundances, and mineral compositions of lunar samples, material from key localities representing the early Earth, and meteorites from Mars and igneous asteroids. Laboratory facilities needed to conduct these types of studies are relatively well established within Australia. New types of mass spectrometers that will provide greater sensitivity and open new lines of enquiry are likely to become available within the decade and should be sought by departments involved in this type of research.

2c. Planetary geodynamics

Planetary geodynamics is the study of interacting natural processes that operate on and within planets. This type of research integrates surface geology with geophysical perspectives of planetary interiors to characterize structures and processes on regional and global scales. The primary datasets are typically derived from spacecraft observations and, where available, landed instruments.

Theme #2: How and why do planetary surfaces, atmospheres, and climates change over time?

2c-1. Surface environments

Key Science Question #2c-1:

* In what way do the early surface environments and interior processes of Earth and Mars resemble each other, and how and why have the subsequently diverged?

Significance of Key Science Question #2c-1:

Early Earth and Mars both appear to share similar dense atmospheres, a methane-CO₂ greenhouse, global magnetic field, a hydrologic cycle, abundant volcanism, and the effects of the LHB. The extent of this similarity and when and how the planets diverged to their present different states may highlight common principles of planetary evolution

Background

Much of the research focus over the coming decade will be directed toward a comparison of the structures and evolution of the terrestrial planets Mars, Venus, Earth and the Moon in order to understand the diversity of planetary bodies within our Solar System, their magmatic and tectonic histories, and their atmospheres and climate systems. The scale of investigations ranges from near-surface processes such as weathering, hydrothermal alteration, and wind, water and ice erosion and deposition that modify and shape crustal structure, composition and surface morphologies of the planets, to megascopic processes such as mantle convection and meteorite impacts.

An important mechanism for the transfer of heat and mass within solid planets and ultimately to the hydrospheres and atmospheres, is the motion of magma. Volcanic and tectonic activity is a consequence of the release of thermal energy. The various mechanisms that planets adopt for releasing heat is a function of planet size, composition, and distance from the sun. While for Earth plate tectonics and mantle plumes represent efficient cooling mechanisms, exploration of Mercury, Venus and Mars suggests that there are other thermal mechanisms that should be subtly reflected on the surface of planets by volcanic and tectonic features. Therefore, it is to be expected that the study of

terrestrial analogues work may be valuable but not sufficient for understanding how other planets work.

In general, the evolution of planetary atmospheres and climate are strongly coupled to global tectonic styles. Much can be gained from learning how the past and current climates on the other terrestrial planets evolved to their present state, from the release of volcanic gasses to the role of biota in the carbon cycle and erosion. Internally generated magnetic fields shelter the surfaces of planets from the solar wind and intense radiation, yet requires recycling of the lithosphere to the core-mantle boundary in order to drive convective currents inside a liquid core strong enough to generate a dynamo. Does this very nature of the continual recycling of the Earth's crust (which exchanges water and gases between the atmosphere and deep interior) provide the materials and energy necessary for developing and sustaining life on the planet?

A key aspect for understanding the evolution of other planets over the coming decade will be exploring their surfaces, and placing what we see in the context of geological processes that shaped the planets. The success of the Mars exploration Rover missions means field geology studies of planetary surfaces can be coupled with remote sensing observations by orbiting spacecraft and the study of spot locations by immobile landers. Field studies of planetary surfaces now have a higher research and public profile since the end of the US Apollo and the Soviet Lunakhod missions in the 1970's. This focus will continue with the short-term prospect of a flotilla of orbital spacecraft and rovers to Mars (e.g., MSL in 2009, ExoMars in 2011) and the Moon by several nations including China, Japan, and India as well as Europe and the USA. The longer term prospects grow more exciting with landed missions to the Moon expected within the next 8-10 years and the resumption of human exploration of space beyond low-Earth orbit within the next 15-20 years.

Australia has various research groups that have been very productive in recent years on the specific topic of numerical modeling of mantle convection on a planetary scale. Australia also has substantial expertise in remote sensing and mapping, important in defining and exploring relevant surface features of planets. The combination of these two major lines of investigation could provide improved thermal models of the terrestrial planets of the solar system. This has even broader repercussions, also to complement cosmochemical studies aimed at constraining the detailed composition of the interior of planets.

Experiments and facilities needed to address Key Science Question #2c-1:

Characterisation of the stratigraphy and composition of the Martian crustal units at the regional and (where possible with lander and rover data) outcrop scales, and developing of testable hypotheses for their formation. This is to be achieved by (1) harvesting and processing of remote sensing data available from previous and ongoing missions to Mars, as well as (2) by direct involvement of Australian scientists in the planning stages of future missions to Mars aimed at collecting specific data to address the question.

(1) There is a need in Australia to establish a hub of planetary data harvesting and processing activities, supported by computer infrastructure, data storage capacity, and personnel straddling modelling, computer processing, physical and geological expertise. A core of such activities already exists at the University of technology, Sydney, where the procedures to process data from the Mars Odyssey THEMIS (Thermal Emission Imaging System) and Mars Global Surveyor MOC (Mars Orbiter Camera) instruments and to analyse the results using a geographic information system (GIS) are already routinely implemented in studies of the surface of Mars. Additional support in this area would allow the set up of a facility to establish a formal NASA Planetary Data Node in Australia, to increase the number of image processing tools (for example GMT and VICAR) and make them available to other groups in Australia in a user-friendly format that facilitates examination of the data as well as providing expertise to model, process and interpret data.

(2) Several regions of Mars have been investigated at a scale that has been useful to constrain general tenants of Mars genesis and evolution, but that has opened new questions to be addressed by studying features at different scales. This requires design and production of experiments and equipment to be deployed in future missions to the planet. However, the lack of formal collaborative agreements between Australia and the space-faring countries means that Australian scientists are effectively precluded from participation as principal investigators able to propose experimental designs and produce and test prototypes and final equipment to participate in future missions. Establishment of formal international collaborations between Australia and the agencies operating future planetary science and exploration missions is indispensable to broaden Australian participation in the study of Mars and other planetary bodies. This will produce two outcomes: firstly, Australian scientists will have priority for the examination and analysis of the data; secondly, Australian industry will be directly involved in the design and production of equipment. This will generate know-how in Australia and create the conditions for technological development in areas in which Australia can become a world leader and unique provider.

2c-2. Planetary atmospheres

Key Science Question #2c-2:

* Why have the atmospheres of Earth, Mars, Venus, and Titan followed very different evolutionary paths?

* How has greenhouse warming kept the Earth habitable, but not the other planets?

Significance of Key Science Question #2c-2:

Planetary atmospheres are important for the ways they shape planetary surfaces through erosion and for the controlling influence they have on climate. On Earth, the atmosphere is critically linked to the biosphere because all the important atmospheric gases, with the

sole exception of argon, are biologically mediated to some extent. Understanding the evolution of a planet's atmosphere, therefore, provides an important perspective on the history and habitability of a planet.

Background

There are four terrestrial bodies in the Solar System with significant long-lived atmospheres: Venus, Earth, Mars, and Titan. However, each of these atmospheres have distinctive characteristics, reflecting the unique evolution of the planet. The atmospheres of Venus and Mars are mostly carbon dioxide, while those of Earth and Titan are mostly nitrogen. Atmospheric pressures also vary widely, from ~90 bar on Venus, to 1 bar on Earth, and 0.07 bar on Mars, despite probably similar starting compositions.

The atmosphere of Venus is very acidic with high concentrations of hydrochloric and hydrofluoric acids, and sulfur dioxide that readily reacts with other gases to make sulfuric acid. Earth's atmosphere is predominantly nitrogen but it is distinguished by abundant oxygen, reflecting the presence of photosynthetic life. Titan's air nitrogen atmosphere on the other hand is unique for the presence of reduced hydrocarbons. Factors that contributed to the formation and modification of planetary atmospheres included primary contributions from trapped nebular gas, erosion by solar wind, thermal escape, impact, atmospheric cratering, condensation, and chemical reactions.

Experiments and facilities needed to address Key Science Question #2c-2:

The surface and lower atmosphere of Venus is largely hidden from direct view by the dense sulphuric acid clouds that extend up to 70km altitude. During the 1980s Australian astronomer David Allen, using the Anglo-Australian Telescope at Siding Spring, discovered a means of seeing through these clouds by observing the nightside of the planet at near-infrared wavelengths. Through "windows" at certain infrared wavelengths it is possible to see thermal radiation from the lower atmosphere, and surface. This makes it possible to study the atmospheric composition and properties in regions that would be hard to reach using in-situ probes because of the extreme temperatures and pressures.

This technique, developed in Australia, is being exploited by the ESA Venus Express spacecraft, now in orbit around Venus. Australian space scientists are also using this technique for continued studies of Venus from the Anglo-Australian Telescope using its new infrared spectrometer IRIS2. Some of the latest results from these ground-based observations of Venus show how infrared observations can be used to probe different levels of the Venus atmosphere, and how they can complement the data obtained by Venus Express. These data will be used to study the composition of the atmosphere near the surface, to study the composition and circulation of the cloud layers, and to follow the highly variable oxygen airglow emission, which provides a probe of upper atmosphere chemistry and dynamics.

2c-3. Australian analogues for planetary surfaces

Key Science Question #2c-3:

* How can studies of Australian geological processes (magmatic, sedimentary, hydrothermal, and geomorphological) better inform our understanding of the other terrestrial planets?

Significance of Key Science Question #2c-3:

The extension of an intelligent uniformitarianism to other planets requires and understanding to the degree to which common and unique processes exist between them. Terrestrial examples can serve both as analogues for extraterrestrial processes, and also as yardsticks against which differences resulting from different compositions, thermal and tectonic regimes, or planetary histories, can be measured.

Background

In addition to theoretical studies of global scale physical and chemical processes, there is an opportunity to use Australian environments as natural laboratories to better understand extraterrestrial surfaces, test hypotheses, and develop criteria for recognition of specific processes. For example, the epicratonic basins, landforms and regolith of the Australian interior, such as that near Arkaroola, provide many analogues to the Martian landscape. The longitudinal dune systems appear to have close parallels on Titan, despite the different atmospheric density, surface composition, and temperatures. Australian researchers are world leaders in the understanding of these systems. The Australian continent has a diverse record of impact cratering and the modification of the craters through hydrothermal alternation, burial, and exhumation. Many of these impact structures have received only cursory study. Finally, Australian geologists have unparalleled expertise in working with early crustal rocks in the Pilbara and Yilgarn Cratons. We suggest that studies of such Australian analogues be a priority within the space science decadal plan.

The topic offers at least three major opportunities: 1) development of new or improved instruments for the collection of *in situ* data (using Australian expertise in spectroscopy, microbeam techniques). 2) Field trials of individual instruments or full-scale prototype rovers in terrestrial analogues settings (such as the FIDO rover trials in the SW United States or the NOMAD trials in Chile and Antarctica). 3) Framing and testing of hypotheses for field-scale extraterrestrial features with reference to their terrestrial counterparts, especially those features which are particularly well developed in Australia (the results of acid-groundwater systems, stable ancient landscape processes, small impact craters, sief dunes).

Experiments and facilities needed to address Key Science Question #2c-3:

Better development of analogue regions in Australia, e.g. the Pilbara, Arkaroola, various impact craters, and other locations, as required.

2d. The Impact of Impacts

One of the enduring legacies of planetary science is an appreciation of the importance of large-scale collisions or impacts as a fundamental process, especially during the early stages of planetary evolution. An accurate reading of this impact history is important for establishing the significance of large impact events for crust formation and biologic evolution on Earth, absolute timescales of geological events on other planets, and planetary dynamics in the Solar System.

Theme #3: What is the role of meteorite and asteroid impacts in planetary evolution?

Key Science Question #2d-1:

- * Was there a cataclysmic bombardment of the inner solar system about 3.9 billion years ago and if so, where did the impactors come from?
- * Is there a signature of the late heavy bombardment in the oldest Australian rocks and minerals?

Significance of Key Science Question #2d-1:

Resolving this question would answer a long-standing problem in planetary science and would address current controversies over the source of impacting planetesimal populations that created the large nearside lunar basins. Numerical modelling has raised the possibility that the outer planets Neptune and Uranus either formed late or migrated away from the sun ~500-700 million years after formation of the terrestrial planets. This may have stirred primitive, icy objects from the Kuiper Belt, sending them crashing toward the inner solar system. In contrast, the size distribution of lunar craters is more consistent with a provenance for the impactors in the inner solar system, probably asteroid belt. Distinguishing among these alternatives would provide a better understanding of the evolution of the solar system after the primary planetary structure was established.

Background

A major insight gained from the study of lunar samples was the realisation that massive impact events occurred considerably later than most models of planetary accretion would have predicted. The age distribution of lunar impact melt rocks and glasses show that a population of large (10-100 km diameter) planetesimals struck the lunar surface at ~3.8-4.0 Ga, some 500-700 million years after initial differentiation of the crust and mantle. The heavily cratered surfaces of Mars and the ages of meteorites from the asteroid belt suggest that a late bombardment at ~3.9 Ga was a general feature of the inner solar system. Where these impactors came from, why they invaded the inner Solar System at this particular time, and their possible influence on the evolution of the terrestrial planets are hot topics in planetary science.

The late heavy bombardment occurred at a time just before the earliest evidence for life on Earth and has important implications for the origin and early evolution of life. The cataclysm theory leads to much lower total impact rates on the very early Earth than the “steady decline” model. This opens up the possibility that life on Earth could have started very early, and survived through the cataclysm. Alternatively late bombardment may itself have played a role in the origin of life, through delivery of organics to the Earth, creation of temporary environments, or transferring material between planets.

Experiments and facilities needed to address Key Science Question #2d-1:

Studies that establish a high-resolution record of the timing of impact events and the provenance of planetesimals traversing the inner solar system are needed. The chronology of early impact events in the inner solar system and the provenance of planetesimals that bombarded the Earth and Moon can be established through geochemical studies of the ages and chemical compositions of lunar impact melt breccias and glasses. The age and composition of the earliest terrestrial crustal rocks and minerals need to be studied to identify whether they contain a signal of the impact record on the early Earth.

Key Science Question #2d-2:

* What role did impacts play in the long-term evolution of the Australian continent and surface environments?

Significance of Key Science Question #2d-2:

Improved understanding of the magnitude, rate, and location of impacts onto the Australian continent over the 3.7 billion years of its history will provide better understanding of one particular aspect of the origin and evolution of the early Earth. This is of significance to understanding the formation the compositionally unique terrestrial crust, constraints and niches for Earth’s earliest biosphere, and for the development of Archaean metallogenic processes.

Impact rates fell away dramatically after ~3.9 Ga but were still significant. The Archaean (2.5-3.7 billion years old) rocks of the Pilbara contain numerous horizons rich in spherulitic glasses that demonstrate major impact events continued to hit the Earth. These spherule horizons have been used to provide tentative correlations between the Pilbara and Southern African cratons. Although no Archaean impact sites have been identified, the chemistry of the spherules provide constraints on the possible target rocks, which may include lithologies, such as Archaean oceanic crust, no longer present on the Earth’s surface. Furthermore, the rate and magnitude of impact events may have constrained the development of the earliest terrestrial organisms.

Experiments and facilities needed to address Key Science Question #2d-2:

Detailed geochemical and isotopic studies of Archaean spherule horizons, search for new localities in Australia and overseas.

Key Science Question #2d-3:

* What has been the role of impacts in biospheric evolution?

Significance of Key Science Question #2d-3:

Large impacts have the potential for significant disruption of the biosphere but the links between major impact events, biosphere disruption, and mass extinction are poorly understood. The best known of such events is the Chicxulub impact, which played a key role in the Cretaceous-Tertiary mass extinction. Other extinctions have been linked to impact events, although links for these are not as clear as for the K-T event.

Better understanding of the processes occurring in the atmosphere and on the ground during a large impact event would greatly improve understanding of the evolution of the biosphere, the history of biodiversity, and the risk posed by large, rare events. Two events in particular stand out for further investigation, the Acraman impact event (590 Ma) and the Woodleigh and Picaninny structures (~360 Ma) in South Australia and Western Australia, respectively. Better understanding of the magnitude and frequency of impact events through the Phanerozoic would also constrain the orbital dynamics of Earth-cross comets and asteroids.

Experiments and facilities needed to address Key Science Question #2d-3:

Detailed stratigraphic, sedimentologic, palaeontological, and geochemical investigations of outcrops and drill holes in Australia and overseas across the relevant stratigraphic intervals in the Ediacaran and Devonian periods. Modelling of the environmental consequences of the known impacts against the magnitude of the extinction events would be valuable.

Key Science Question #2d-4:

* What role do impact structures play in modifying groundwater and hydrocarbon flow paths and prospectivity in Australian Phanerozoic sedimentary basins?

Significance of Key Science Question #2d-4:

The groundwater and hydrocarbon flow paths and prospectivity in Australian Phanerozoic sedimentary basins is of critical importance to present and future hydrocarbon exploration in Australia and to selection of sites for CO₂ sequestration. A number of medium (4 km) to large (120 km) impact structures are known from Australian Phanerozoic sedimentary basins. Examples include Bedout, Talundilly, Woodleigh, Yallalie, Gosses Bluff, Tookoonooka, and Mt Toodinna. Some of these are in basins with known petroleum prospectivity. The impact structures modify fluid flow paths in rocks surrounding the impact site to a distance of at least one diameter away from the impact rim. Changes to porosity, permeability, and subsurface structural and to the

thermal history of the target successions can potentially alter petroleum source rocks, reservoirs, and trap potential. However most of these impact sites have been very poorly studied to date.

The consequences of large impacts into mechanically weak and/or water saturated sediments is poorly constrained by field studies of terrestrial examples. A better understanding such impacts is important to improved models of the environmental consequences of terrestrial impacts. It is also important for improved understanding of impact processes on Mars, where many impacts appear to have struck targets composed of sedimentary, water saturated material.

Experiments and facilities needed to address Key Science Question #2d-4:

Significantly improved 3D geological and geophysical characterisation of Australian impact craters in basinal successions coupled with improved modelling of impact dynamics and comparison with possible counterparts on Mars.

Key Science Question #2d-5:

* What is the impact risk to Australia?

Significance of Key Science Question #2d-5:

The urbanized, networked and industrialized societies of the 21st century are vulnerable to even small impact events. These include small (Hiroshima-size) to medium (Tunguska-size) air-bursts that occur with once a year to once a century, frequencies, respectively, to direct consequences of a land impact, to the indirect consequences such as tsunamis of an ocean impact. There is a significant hazard of small earth-cross bodies causing local to regional scale devastation. Current knowledge of the risk is very limited and early warning minimal. Unlike other natural hazards, the impact risk is one that can be, at least in principle, minimised by direct action. Furthermore, quantification of the risk posed by such events will generate significant data on the composition and orbits of small bodies in near Earth space, with applications to a wide range of astronomical and planetary science questions.

Experiments and facilities needed to address Key Science Question #2d-5:

Establishment and/or upgrading of long term monitoring ground-based networks of optical and radar systems to track near earth objects. These may be supplemented by microsatellite systems in earth orbit.

2e. Habitable Planets and the Uniqueness of Earth

Of all the questions potentially answerable by planetary science, “Are We Alone” may resonate most deeply with the public. Phrased somewhat less existentially, answering the question of whether life exists elsewhere in the universe is the ultimate goal of astrobiology.

Theme #4: Is there life elsewhere in the solar system or beyond?

2e-1. Origin of Life

Key Science Question #2e-1:

- * Can we sharply define targets for the exploration for life or former life elsewhere in the Solar System?
- * Do early habitats of terrestrial life indicate unique tectonic or chemical environments are needed for the origin of life?

Significance of Key Science Question #2e-1:

The search for life and intelligence beyond Earth is of profound scientific and cultural significance. Knowing whether or not the Earth is representative of a much larger population of habitable worlds beyond the Solar System is a key part of this search.

Background

Astrobiology is the study of the origins, evolution, distribution, and future of life in the universe. This broad field embraces the search for potentially inhabited planets within and beyond our Solar System, including the exploration of Mars and the outer planets, laboratory and field investigations of early life on Earth, and studies of the potential of life to adapt to future challenges, both on Earth and in space. Astrobiology is interdisciplinary, combining planetary science, astronomy, and space exploration technologies with molecular biology, ecology, information science, and related disciplines. The broad interdisciplinary character of astrobiology requires a comprehensive and inclusive understanding of biological, planetary and cosmic phenomena.

Experiments and facilities needed to address Key Science Question #2e-1:

Research is needed to understand the mechanisms by which habitable environments have evolved throughout the Solar System and how the planetary environments have influenced the evolution of life. Through study of the reciprocal interactions of organisms and their planetary environment we strive to develop an understanding of the biochemical and metabolic machinery that drives the global physical and chemical cycles. We need to

understand aspects that seem to make Earth particularly well-suited to support complex life, for example the possible significance of tectonic environment and how it has evolved through time, and the role that delivery of complex organics by primitive bodies may have played in setting the initial conditions that allowed life to begin.

2e-2. Search for Earth-like planets

Key Science Question #2e-2:

- How special is the Earth?
- Are there Earth-like planets elsewhere in the galaxy?

Significance of Key Science Question #5-1:

Studies of extrasolar planets¹ (exoplanets) over the next decade will help science address the fundamental problem of whether the Solar System and our own planet have close analogues elsewhere in the Galaxy.

Background

In observations that complement more detailed Solar System analyses, astronomical surveys using different techniques are revealing some 170 gas giants and what appear to be rocky planets in orbit around other stars^{2,3,4}. The majority of extrasolar planets are found using stellar radial velocity measurements, the so-called "Doppler wobble" technique. These ground-based radial velocity surveys use telescopes equipped with a precision spectrographs to measure the subtle wavelength shift in a star's light due to the motion of the star and its planet around their common centre of mass. The Doppler wobble technique is the main method used to detect planets around "nearby" stars within about 100 light years from us.

A transit method also can be used to find exoplanets, typically by simultaneously monitoring a field of huge numbers (many thousands) of distant stars. Detection of a planet involves recording the temporary dimming of a star's light by a planet whose orbit happens to make it transit in front of the star during the monitoring period.

Gravitational microlensing forms the basis for another exoplanet detection technique for distant stars. If a star and its attendant planet passes between us and a more distant background star, the light from the background star is focused towards us by the gravity of the foreground star and its planet. This effect briefly brightens the background star in a characteristic way that reveals the planet's presence.

¹ http://en.wikipedia.org/wiki/Extrasolar_planet

² <http://planetquest.jpl.nasa.gov/index.cfm>

³ <http://exoplanets.org/>

⁴ <http://www.dtm.ciw.edu/boss/planets.html>

The direct imaging of planets is also now being pioneered.

The surprising overall result from the first decade of exoplanet observations 1995-2005 is the apparent rarity of planetary systems similar to our Solar System. Prior to 1995 we expected to find Solar System analogues with gas-giant Jovian planets in distant orbits, and inner terrestrial planets, all in near-circular orbits. Instead, almost every exoplanet found to date turn out to be gas-giant Jovians in highly eccentric orbits or "hot Jupiters" in extremely close, tidally-circularised orbits. To make matters worse, present information on terrestrial exoplanets is almost non-existent, due to the difficulty of detecting such small, faint and low-mass worlds.

Despite advances made over the past decade, to what extent the Solar System is unique remains very much an open question. This is because observations of extrasolar planets have not been operating long enough to detect gas giant planets in the distant long-period orbits similar to Jupiter and beyond, and as yet, we have little credible information on terrestrial exoplanets.

Over the next decade, observational advances for the first time will enable meaningful comparisons of our Solar System with other planetary systems. Australia currently hosts some active extrasolar planet search programs and the nation has the potential to further develop its current expertise and scientific successes in detecting and characterising exoplanets over the next decade. Australia's participation in exoplanet science covers different observational techniques, as well as some theoretical studies. Radial-velocity measurements are made for the Anglo-Australian Planet Search using the Anglo-Australian Telescope.⁵ Transit and microlensing-based searches are in operation (such as the PLANET collaboration), or are in development. These searches are based on infrastructure in place or funded for Siding Spring Observatory in NSW and Mt Canopus Observatory in Tasmania.^{6,7,8} Innovative development work is also being done on an instrument for the direct detection of exoplanets using polarimetry.⁹ Theoretical modelling studies are conducted at several of Australia's universities.^{10,11,12}

Over the next decade, progress in Australian exoplanet science will include more ground-based radial velocity survey that will achieve a long enough time base of observations to readily detect a host of Jupiter-like worlds in Jupiter-like orbits around nearby stars (as long as these planets actually do exist). In addition, developments in hardware and software in radial velocity measurements and transit and microlensing efforts should lead to a more comprehensive set of discoveries of terrestrial worlds orbiting nearby and distant stars. These ground-based efforts also pave the way for more detailed space-based

⁵ <http://www.aao.gov.au/local/www/cgt/planet/aat.html>

⁶ http://arxiv.org/PS_cache/astro-ph/pdf/0211/0211098.pdf

⁷ <http://www.physorg.com/news10228.html>

⁸ <http://www.phys.unsw.edu.au/astro/research/thesis/MartonHidas.pdf>

⁹ <http://aca.mq.edu.au/People/Bailey.htm>

¹⁰ <http://www.mso.anu.edu.au/PSI/>

¹¹ <http://hubblesite.org/newscenter/newdesk/archive/releases/2005/10/video/b>

¹² <http://online.itp.ucsb.edu/online/astro99/mardling/>

studies, and provide the observations on which theoretical advances can be made in understanding the formation and evolution of planetary systems. Solid evidence also should emerge as to the frequency of different types of planets and planetary system architectures. This will help us understand how widespread are planetary systems like the Solar System, and how unique or commonplace are habitable terrestrials like the Earth.

Can we sharply define targets for the exploration for life or former life elsewhere in the Solar System? Although phrased in very general terms this question encompasses many aspects of planetary science but at the same time has a clear goal. It leads to emphases on Mars, Europa, Io, Enceladus and Titan but in one way or another encompasses all objects in the Solar System, the history of the energy output from the Sun, the history of the Earth-Moon system, and more. It links us clearly into the exploration programs of NASA and ESA. It includes instrument development.

Experiments and facilities needed to address Key Science Question #2e-2:

Ground-based planet-searching programs in Australia need continuing financial support for their maintenance and development, to garner maximum scientific results from established and developing infrastructure at Siding Spring Observatory and Mt Canopus Observatory. Of particular note is the need to persist with the radial velocity work with the Anglo-Australian Telescope, as the data from this program dates back to 1998, and so offers a unique set of long-term measurements for detecting Jupiter analogues around nearby stars in the southern sky.

3. Summary of Australian planetary geoscience research concentrations

The following submission was presented by the Geological Society of Australia Specialist Group in planetary science to the NCSS strawman solicitation in October 2005. It identifies areas of existing national strengths in planetary geosciences.

GEOLOGICAL SOCIETY OF AUSTRALIA Incorporated

Registered Office

Suite 706

301 George Street

SYDNEY NSW 2000

AUSTRALIA

Phone: (02) 9290 2194

Fax: (02) 9290 2198

Web: www.gsa.org.au

ABN: 93 652 757 443



Dr Iver Cairns
Chair, National Committee for Space Science
School of Physics
University of Sydney NSW 2006

20 October 2005

Submission to the Australian Decadal Plan for Space Science

Dear Dr. Cairns:

Planetary science will be one of the most exciting and rapidly expanding research areas of the coming decade. The recent discovery of more than 130 extrasolar planets and the continuing exploration of our Solar System through spacecraft missions and studies of extra-terrestrial materials provides powerful stimuli for understanding the geological and astrophysical processes that create planetary systems and shape the surfaces and interiors of planetary bodies.

Planetary science is inter-disciplinary and international in scope, linking earth scientists, astronomers, physicists, chemists, biologists, engineers, and mathematicians working independently on topics as diverse as the origin of our Solar System, global climate change, and development of new technologies for laboratory analysis and remote sensing. The knowledge base among planetary scientists in Australia is vast but somewhat disjointed and less effective than it might be, mainly because of lack of coordination at the national level. National support will drive breakthrough science, grow new technologies, and enhance Australia's international prestige.

This submission is from the Specialist Group in Planetary Geoscience, which was created under the auspices of the Geological Society of Australia with the following goals:

- Provide a national focus for research into the origin and evolution of planets and planetary systems.
- Support Australian research on extra-terrestrial materials, the origin of the solar system, surface processes on other planets, the nature of planetary interiors, the

effects of impacts, and the long-term development and evolution of planetary atmospheres and climatic systems.

- Advance national strategic goals for research and education in geosciences relevant to planetary studies.
- Encourage development of national and international linkages with other societies and organisations that benefit Australian geoscience.
- Promote planetary geoscience to the broader community.

The GSA-SGPG represents planetary geoscientists from over 25 different educational, governmental, and industrial institutions across Australia. The development of an Australian Decadal Plan for Space Science is most welcome, and the GSA-SGPG thanks the Academy's Space Science national committee for the opportunity to present this submission.

The GSA-SGPG has identified the following areas of existing national strengths in planetary sciences:

*** Geophysics of Stellar Disks and Planetary Interiors**

This work concentrates on modeling the 3D thermal, chemical, and mechanical evolution of stellar disks, planetary orbits, and the interiors of planetary bodies on scales ranging from crustal thickness to the entire globe. Australian research groups have developed an extensive repertoire of modeling algorithms and implementations for handling the strongly non-linear, history dependent constitutive properties of planetary materials, and dealing with the large range of emergent length scales. The primary Australian research groups involved in modeling solar system dynamics and planetary geophysics are at Monash and the Australian National University, with nuclei developing at the University of Queensland and Sydney University.

*** Astrobiology and Extra-solar Planets**

The question of how life began on Earth, and possibly elsewhere in the solar system and universe, is one for which Australia holds unique natural resources, such as the oldest known minerals and fossils. However, the leap from abiotic organic precursor molecules to functioning metabolic processes remains mysterious. Addressing this issue will involve understanding the co-evolution of the Earth and its biosphere, issues such as the history of life's origin, timing, and critical intervals, and the interaction of extraterrestrial, geological, and biological processes.

The study of planetary systems around other stars is a rapidly developing, astronomically-based field that provides an observational basis for evaluating the diversity of planetary habitats across the galaxy and the universe. Significant national strengths in astrobiology and extra-solar planetary systems have been developed at Macquarie University, The University of Western Australia, and The Australian National University.

* **Cosmochemistry**

Understanding how the Solar System formed and evolved requires knowing about the chemistry of planets and their precursors, and the ways in which elements are created in stars. Cosmochemistry is the laboratory study of extraterrestrial materials (meteorites, cosmic dust, lunar samples) aimed at understanding the formation and development of the Solar System from a chemical perspective.

There are significant national strengths in cosmochemistry in Australia, including groups at the Australian National University, Monash University, Curtin University, Macquarie University, and the Western Australian Museum. Current cosmochemical research in Australia is focused on the time scales of planetary formation, processes that construct and modify planets on global scales, conditions in the solar nebula, and the delivery of supernovae and interstellar products to the early solar system.

The coming decade will present significant opportunities for discovery as spacecraft missions to asteroids, comets, and the Moon return new types of extra-terrestrial material to Earth, new meteorites continue to be discovered in Antarctica and the hot deserts of the world, and new analytical facilities are developed.

* **Australian Analogues of Mars Environments**

The ability of planetary geoscientists to extrapolate their understanding of physical processes that have shaped the Earth to other planets provides the most fundamental and advanced interpretative tool for building comprehensive models of the evolution of Mars and martian environments. Analogue studies related to aeolian, flood, periglacial and glacial, marine, volcanic, tectonic and impact processes bear directly on our understanding of climatic, hydrologic, and geologic conditions on Mars and its potential for life. The exploration of Mars has captured the imagination of the public, providing a vehicle for education on a variety of scientific issues relevant to planetary environments.

Australia represents one of the most versatile grounds in the world to conduct analogue studies of the martian environment, and there is broad national interest in this type of research, which links space science, earth science, and technology development. Research activities are diverse and range from spectral mapping of alteration mineralogy using airborne instruments for predicting signatures of martian hydrothermal systems, to modelling the geography of planetary surfaces using spatial analysis of landform distributions, patterns and relationships between comparative Mars-Earth land systems. The 2004 meeting of the Australian Geological Convention hosted special sessions on martian analogue research, emphasising Australian connections and contributions. This was followed by a thematic issue of the Australian Journal of Earth Sciences (June 2005) highlighting Australian-led research on Mars analogues in the Australian geological record.

This type of research has the potential to generate national benefits in the areas of

information and instrumentation technology, as well as scientific gains. For example, the GeoSpatial Group at the University of South Australia aims to develop a geographical information and analytical system for planetary data. Research by CSIRO and commercial companies have delivered a suite of instruments that allows spectral mineral mapping and analysis at a range of scales. These systems have been widely used in mineral exploration and regolith research in Australia and overseas, and they provide a strong technological base for the development of spectral instruments for planetary exploration by robot rovers, automated drilling systems, or flyby and orbiter probes.

An important infrastructure component for Mars analogue research is the establishment of a field station in the Northern Flinders Ranges, where operational and laboratory tests can be conducted. These capabilities will support planning and exploratory missions to Mars. While still in its infancy, this initiative, if adequately supported in Australia, has the potential to become an essential asset for mainstream international space missions to the red planet, and as such it has attracted attention from both NASA and the European Space Agency. The availability of a large planetary science knowledge base in Australia makes this initiative particularly attractive to potential international collaborators.

* **Meteorite impacts – the Australian record**

The old age and stable surfaces of Australia provide one of the best regions worldwide for exploration of the impact history of Earth. There are 26 established impact structures in Australia, and at least that many more structures for which an origin by meteorite impact has been proposed. The record of impact events in Australia spans 3.5 billion years of earth history, from the oldest known impact ejecta in the Pilbara to young, well-preserved craters such as Wolfe Creek and Woodleigh, the world's 4th largest impact structure. Over the last few years a small group of dedicated geoscientists have established a discovery rate of approximately one impact structure per year, using sophisticated geophysical and geochemical methods.

Current research programs include examination of potential relations between large impacts, large volume volcanic eruptions, and global scale environmental changes. To date most impact studies in Australia have been conducted on an individual basis with scant funding. As an indication of the level of national interest in this topic, the Australian Journal of Earth Sciences recently published a volume in honor of Eugene Shoemaker that contains 22 papers updating the Australian impact record, and placing it within the context of planetary and space sciences.

* **Spacecraft missions**

Individual Australian geoscientists in various institutions across the country are involved in NASA and ESA space missions, albeit typically at a secondary level. A number of Australian geoscientists have collaborations with the International Research School of Planetary Sciences in Pescara, Italy, and either are negotiating for or have

been granted direct access to data from the ESA Mars Express mission High Resolution Stereo Camera and the Mars Advanced Radar for Subsurface and Ionosphere Sounding experiments.

Considerable effort in Europe, Japan, and the USA is being directed toward sample-return and remote sensing missions to the Moon, Mars, and small bodies such as comets and asteroids. Australian geoscientists are well positioned to participate in these missions, a fact clearly recognised among colleagues outside of Australia as the individual collaborations attest. For example, Australian cosmochemists have been invited by the Japanese MUSES-C asteroid sample return mission to serve on the preliminary examination team and conduct invited research on the returned material.

The scale and scope of Australian involvement in spacecraft missions could be magnified through national support for space science. For example, formal links between Australia, European countries, Japan, and the USA to allow Australian scientists direct access to mission planning and data retrieval could be encouraged and actively pursued. Support for mission participation by Australian scientists would benefit the nation by providing more direct access to technology developments, in addition to a higher profile for Australian scientific achievements.

Sincerely,

Dr. Marc Norman
Australian National University

Dr. Graziella Caprarelli
University of Technology Sydney

Dr. Louis Moresi
Monash University

Prof. Malcolm Walter
Macquarie University

Dr. Jon Clarke
Geoscience Australia

Dr. Andrew Glikson
Australian National University

On behalf of the Geological Society of Australia Specialist Group in Planetary Geoscience.

NCSS Decadal Plan - Planetary Sciences Working Group

Contribution by Dr. Jeremy Bailey

Planetary Astronomy Group
Australian Centre for Astrobiology (ACA)
Macquarie University, Sydney
June 27 2006

Our Interests

The ACA planetary science research involves observation and modelling of planetary atmospheres and surfaces. We use, in particular, near-infrared spectroscopic imaging techniques, and polarimetry to observe planets in the solar system as well as extrasolar planets. We are specifically working on the atmospheres of Venus, Mars and Titan. To interpret remote sensing observations of planetary atmospheres we have developed a new planetary atmosphere radiative transfer model (VSTAR – Versatile Software for Transfer of Atmospheric Radiation). This is a high-spectral-resolution model incorporating full multiple-scattering radiative transfer solutions.

Involvement in Space Missions and Other Projects

ExoMars — We proposed a concept for an infrared spectrometer to go on the ESA ExoMars rover. The spectrometer would be used to detect minerals indicative of promising locations for past life.

Deep Impact — We participated in the Earth-based observing program to observe the encounter of the NASA Deep Impact spacecraft with comet Tempel 1. We helped to organise the Australia-wide involvement in Deep Impact observations through a workshop held at Macquarie in September 2004.

Venus Express — In collaboration with Frank Mills (ANU), I submitted a successful “Supporting Investigator” proposal for involvement in this ESA mission. This means we will have access to mission data and will participate in the project through coordinated ground-based observations and atmospheric modelling.

Virtual Planetary Laboratory — We are international participants in this NASA Astrobiology Institute team that is developing models for terrestrial planets with the aim of predicting biosignatures that might be observable in extrasolar terrestrial planets.

Martian Meteorology — We are developing techniques for mapping Martian meteorological properties such as surface atmospheric pressure using remote-sensing techniques. At present these methods are being developed using ground-based observations of Mars, but they could be incorporated in a future Mars orbiter mission to provide a Martian weather monitoring and forecasting system.

Future Directions

Clearly we have an interest in future solar-system exploration missions including both orbiters and landers to the Moon Mars and other planets. We are particularly interested in using IR spectroscopy techniques for remote sensing of both planetary atmospheres and surfaces.

Such missions will usually be led by the larger space agencies such as NASA and ESA. This means it is unlikely that Australia will be able to take the lead on any specific project. Rather we need to be able to respond to opportunities as they arise.

There are two issues here. First we need to have something significant to bring to the table in any international collaboration – something that will make Australia attractive as a partner. This means we need to build up expertise in appropriate areas. Two we are working on at ACA are:

1. Ground-based observations of planets — We are currently obtaining some of the best ground-based IR observations of solar-system planets. This can contribute to space missions by allowing easy testing of remote sensing techniques prior to their use in a space mission, by providing long-term studies of planets to help provide context for space observations, and by providing coordinated ground-based observations to support space missions.
2. Radiative transfer models for planetary atmospheres — The VSTAR model currently under development has the potential to provide a relatively easy to use tool for modelling the atmospheres of a variety of planets. Such models are essential to quantitative analysis of most remote-sensing data.

The second issue is that appropriate mechanisms need to be in place to allow Australia to participate in international collaborations with other space agencies. At present it is hard to do this. One problem is the lack of appropriate communication channels with other space agencies due to the lack of an Australian space agency. International collaboration at a substantial level on any space mission is normally on the basis of inter-agency agreements, and that means we are currently locked out of any such involvement.

There are sometimes announcements of opportunity for mission involvement that are open to international participants. However, this normally involves the international participant having to demonstrate that they have funding from their own country to support their involvement. The short timescales for these AOs normally means there is no way that funding could be obtained through methods such as ARC competitive grants. This limits involvement to relatively low levels. A possible solution would be a new funding scheme that specifically supported participations in NASA AOs and similar international programs. It would need to provide rapid decisions, and funding would be conditional on success of the international proposal.