

An 84- μ G Magnetic Field in a Galaxy at $Z=0.692$?

Arthur M. Wolfe¹, Regina A. Jorgenson¹, Timothy Robishaw²,
Carl Heiles², and Jason X. Prochaska³

¹Dept. of Physics and Center for Astrophysics and Space Sciences
University of California, San Diego
La Jolla, CA 92093-0424, USA
email: awolfe@ucsd.edu, regina@physics.ucsd.edu

²Astronomy Department, University of California
Berkeley, CA 94720-3411, USA
email: robishaw@astro.berkeley.edu, heiles@astro.berkeley.edu

³UCO-Lick Observatory, University of California, Santa Cruz
Santa Cruz, CA 95464, USA
email: xavier@ucolick.org

1. Abstract

The magnetic field pervading our Galaxy is a crucial constituent of the interstellar medium: it mediates the dynamics of interstellar clouds, the energy density of cosmic rays, and the formation of stars (Beck 2005). The field associated with *ionized* interstellar gas has been determined through observations of pulsars in our Galaxy. Radio-frequency measurements of pulse dispersion and the rotation of the plane of linear polarization, i.e., Faraday rotation, yield an average value $B \approx 3 \mu\text{G}$ (Han *et al.* 2006). The possible detection of Faraday rotation of linearly polarized photons emitted by high-redshift quasars (Kronberg *et al.* 2008) suggests similar magnetic fields are present in foreground galaxies with redshifts $z > 1$. As Faraday rotation alone, however, determines neither the magnitude nor the redshift of the magnetic field, the strength of galactic magnetic fields at redshifts $z > 0$ remains uncertain.

Here we report a measurement of a magnetic field of $B \approx 84 \mu\text{G}$ in a galaxy at $z = 0.692$, using the same Zeeman-splitting technique that revealed an average value of $B = 6 \mu\text{G}$ in the *neutral* interstellar gas of our Galaxy (Heiles *et al.* 2004). This is unexpected, as the leading theory of magnetic field generation, the mean-field dynamo model, predicts large-scale magnetic fields to be weaker in the past, rather than stronger (Parker 1970).

The full text of this paper was published in *Nature* (Wolfe *et al.* 2008).

References

- Beck, R. 2005, *Lect. Notes. Phys.* 664, 41
Han, J. L., Manchester, R. N., Lyne, A. G., Qiao, G. J., & van Straten, W. 2006, *ApJ* 642, 868
Heiles, C. & Troland, T. H. 2004, *ApJ Sup* 151, 271
Kronberg, P. P., Bernet, M. L., Miniati, F., Lilly, S. J., Short, M. B., & Higdon, D. M. 2008, *ApJ* 676, 70
Parker, E. 1970, *ApJ* 160, 383
Wolfe, A. M., Jorgenson, R. A., Robishaw, T., Heiles, C., & Prochaska, J.X. 2008 *Nature* 455, 638



Birgitta Nordström, Jan Palouš and Hans Zinnecker at Carlsberg.
Ole Strömngren and Aage Bohr in the background.

Photo: Bruce Elmegreen.



Marija Vlajic emphasizing a point during her lecture.