# Quantum Atomic and Nuclear Physics Activities

Activity	used in workshops:	
Band Structure Models	QR7	231
Binding Energy	QI4, QR13	232
Cathode Ray Tube	QR11	233
Classical Particle in an Elastic Potential Energy Well	QR3	234
Colleen's Cubes	QI5, QR14B, QR14T	235
Common Sources of Radiation	QR15B	236
Compact Discs	QR10	237
Coolite Balls	QI4, QR13	238
Diffraction Patterns	QR12B, QR12T	239
Electron Interference	QR2	240
Emission Spectra	QI2, QI3, QR5	241
Exposure Levels	QR15B	242
Hydrogen Spectrum	QI2, QR5	243
Identify the Element	QI2, QI3	244
Laser Light I - Focus	QR9	245
Laser Light II- Spectrum	QR9	246

Light Emitting Diode	QR8	247
Measuring Momentum and Position I	QR4	248
Measuring Momentum and Position II	QR4	249
Measuring Radiation	QI6, QR15T, QR15B	250
Molecular Models	QR6	251
Molecular Models of Semiconductors	QR7	252
Nuclear Power Stations	QR15T	253
Periodic Table	QR6	254
Photoelectric Effect	QR1	255
Potential Wells and Wave Functions	QR3	256
Semiconductor Diode	QR8	257
Smoke Detector	QI5, QR14B, QR14T	258
Solar Panels	QR8	259
Thermocouples and Thermistors	QR8	260
Transistors	QR10	261
Wave and Particle Nature of Light I- Interference Pattern	QI1, QR1, QR2	262
Wave and Particle Nature of Light II- Spectral Lines	QI1, QR1, QR2	263
Waves on a String	QR3	264

X-ray Diffraction	QR12B, QR12T	265
X-ray Pictures	QI6, QR12B, QR12T	266
X-ray Tube	QR11	267

### **Band Structure Models**

#### Apparatus

"egg carton" like containers with indentations for marbles or balls (electrons), divided into bands and spaced appropriately for metals, semiconductors etc, model electrons, "eggs", e.g. marbles, to fill the holes in the containers

#### Action

The students examine the models and explore how electrons and holes can act as charge carriers.

#### The Physics

The "egg cartons" represent energy bands, which can be full, partly full or empty of electrons, depending on the material being represented. More or less electrons can be added to represent n and p type semiconductors.



Accompanying sheet:



Examine the band structure models.

How can you make *n* or *p* type semiconductors?

How can a "hole" act as a charge carrier?

# **Binding Energies**

#### Apparatus

a large chart of the binding energy or mass defect per nucleon

#### Action

The students examine the chart and identify which nuclei are the most stable and which processes can lead to stability for different nuclei.

#### The Physics

The most stable nuclei are those with the greatest binding energy per nucleon. Those that are much heavier than iron tend to undergo fission to move back towards the stable region of the chart. Small nuclei like hydrogen can undergo fusion (at high enough temperatures, such as in stars) to form larger elements with greater binding energy per nucleon.



#### Accompanying sheet:

 Binding Energies

 Examine the chart of binding energies.

 What does the diagram represent?

 Fission and fusion are opposite processes,

 when fission occurs a nucleus breaks apart

 and when fusion occurs two nuclei fuse to form a larger one.

 How can both these processes release energy?

 Which nuclei are more likely to undergo fusion?

 Which will undergo fission? Explain your answer.

232 Workshop Tutorials for Physics – Quantum, Atomic and Nuclear Physics Activities

# **Cathode Ray Tube**

#### Apparatus

working cathode ray tube

#### Action

The students examine the tube and identify the important components. They should note the similarity to the x-ray tube.

#### **The Physics**

The cathode ray tube has a source of electrons (the cathode rays) which are accelerated using an electric potential difference and "steered" using coils. When the electrons collide with phosporus atoms on the screen photons in the visible region are produced. An x-ray tube works in the same way but uses a larger accelerating potential to give the electrons more energy so that x-rays rather than visible photons are released when the target is hit. The target material is usually a metal in an x-ray tube. Old TV sets and oscilloscopes use a phosphorus screen.



#### Accompanying sheet:

### **Cathode Ray Tube**

What is a cathode ray?

The cathode ray tube is very like the x-ray tube.

Identify the main components. How is the cathode ray tube different to the x-ray tube?

### **Classical Particle in an Elastic Potential Energy Well**

#### Apparatus

air track with spring mounted at each end, short glider

#### Action

The students observe the way the "classical particle" (the glider) is reflected from the springs at the ends of the track, and sketch the elastic potential energy of the system as a function of glider position. They should observe that the glider is slowed by the springs and then reflected, and note the relative time spent by the glider around the middle of the track and near the ends. They can then sketch the probability of finding the glider at a given position on the track.

#### **The Physics**

The elastic potential energy, U, of a spring-mass system is proportional to  $(\Delta l)^2$  where  $\Delta l$  is the compression (or extension) of the spring away from its equilibrium length. The spring is compressed by the glider, and is otherwise at equilibrium. The potential energy as a function of glider position, x, is therefore zero except where the glider compresses the springs at either end of the track, at which positions it is proportional to  $(\Delta l)^2$ . This is very much like an infinite potential well, and is approximately square except for the curvature at the edges.

For a given interval in space, the glider spends more time near the ends of the track as it must be slowed down and change direction, then be accelerated away again at the track ends. The surface is approximately frictionless so the glider travels at constant speed once it is no longer in contact with the end springs. The probability density, P, is shown opposite, the dotted lines show the ends of the air track.



The probability density for an electron in a well is the opposite to this for the electron's ground state, and in general is a minimum (zero) at the edges of the well.



# **Colleen's Cubes**

#### Apparatus

100 or more wooden cubes, about 2cm to a side, each with a dot on one side, a large jar and a tray

#### Action

The students pour the cubes from the jar into a single layer in the tray. They remove those with dots showing, count the number remaining and repeat several times, or until no cubes are left. They then sketch the number left with each throw or the number removed. (Or both.)

#### The Physics

This represents the random process of nuclear decay. The sketch will be (approximately) exponentially decreasing.



#### Accompanying sheet:



Shake the jar containing the nuclei (cubes) and pour them into the tray. Write down the number of cubes with dots showing on top, remove those cubes and replace the rest in the jar. How many cubes are left of the original 100? Repeat several times.

Sketch the number of cubes removed (the activity) as a function of time. Sketch the number of cubes remaining in the jar as a function of time.

### **Common Sources of Radiation**

#### Apparatus

some (slightly) radioactive samples and/or laboratory sources, see table below Mantles for gas lanterns are also an excellent example.

#### Action

The students measure the radiation from the sources and compare it to the background level.

#### The Physics

Most of the sources will have levels barely above background. It helps to have a common laboratory source such as Cs for comparison.

Radioactivity in some common substances.

Substance	Activity (Bq/kg)
garden soil	2000
brazil nuts	400
human bodies	80
cows milk	50
sea water	12
tap water	0.1

A student at the Australian Catholic University measuring his own radioactivity.



Dosage limits for ionizing radiation (from the ARPANSA website):

The NHMRC recommended radiation dose limit for the public is 1 mSv (1000  $\mu$ Sv) per year. ANSTO's dose constraint for reactors, which has been agreed by the Nuclear Safety Bureau, is 100  $\mu$ Sv per year for members of the public



### **Compact Discs**

#### Apparatus

CD, reasonably high powered optical microscope

#### Action

The students observe the CD under the microscope.

#### **The Physics**

The students will observe pits, which are spaced to be resolvable by a laser. They should identify that this is how information is encoded on the CD, and the spacing between the pits must be great enough to be resolvable by the CD player. This is digital encoding, unlike magnetic tapes and vinyl records which use analog encoding.



Accompanying sheet:

### **Compact Discs**

Examine the CD under the microscope.

Can you see the pattern of pits which digitally stores the information?

How is this different to the way information was stored on vinyl records? How is this different to the way information is stored on magnetic tape?

# **Coolite Balls**

#### Apparatus

two or three coolite balls hanging from threads very close to each other, some way of charging the balls, e.g. fur and glass rod

#### Action

The students charge up the coolite balls such that they all have the same charge type, and explain their observations. They should make the analogy to putting charged particles together in the nucleus.

#### The Physics

Like charges will repel and the coolite balls will move away from each other. If the only force in the nucleus was the electrostatic force, nuclei would not be stable.



Accompanying sheet:

# Coolite Balls Charge the coolite balls so that they have opposite charges. What happens? Now charge them so they have like charges and observe what happens. How do protons stay together in the nucleus?

### **Diffraction Patterns**

#### Apparatus

laser, piece of pantyhose material in a single layer

#### Action

The students shine the laser light through the material and observe the resulting pattern. They then stretch the material and observe how the pattern changes.

#### **The Physics**

The network of fine threads in the fabric forms a grating. When you shine the laser light through the fabric you see a diffraction pattern.

The spacing between the maxima in the pattern (bright spots) is inversely proportional to the grid spacing;  $d \sin\theta = m\lambda$ , can be used to find the grid spacing, d, given the angular separation,  $\theta$ , of the maxima.

The diagrams show the fabric to the left and the diffraction pattern to the right. When you stretch the fabric horizontally it also squeezes in vertically, the pattern will do the reverse of this, squeezing in horizontally and stretching vertically. When you stretch it vertically it will squeeze in vertically and stretch horizontally.

Note that this activity is also used in a waves workshop.





### **Electron Interference**

#### Apparatus

photographic plates or diagrams showing an interference pattern due to electron interference

#### Action

The students look for evidence of the wave and particle nature of electrons in the pictures.

#### **The Physics**

In the early pictures individual "dots" are visible, where single electrons have been incident on the plate/detector. The discrete nature of the dots shows the particle nature of electrons.

Waves passing through the slits interfere to give a pattern of fringes, with spacing depending on de Broglie wavelength and slit separation. This is visible in the later pictures and shows the wave nature of the electrons.



This diagram, or a similar one, showing electron interference can be used.

#### Accompanying sheet:

### **Electron Interference**

A beam of electrons is directed through two narrowly spaced slits. The emerging beam falls on a sheet of film.

These pictures contain clear evidence that the electrons are behaving like ordinary classical particles (tiny billiard balls).

These pictures also contain clear evidence that that the electrons are behaving like ordinary waves.

### **Emission Spectra**

#### Apparatus

hand held spectrometer, light sources

#### Action

The students observe the spectral lines and compare them for different light sources.

#### **The Physics**

The lines correspond to electron transitions, where the emitted photon has the energy difference between the initial and final electron energy level. This energy and hence the wavelength, can be determined using the Bohr model for hydrogen, but is more complicated for larger atoms. It allows the "fingerprinting" of elements by their unique spectra, a valuable tool for finding out what something is made of.

Optometry students at the University of New South Wales observing a sodium spectrum.



#### Accompanying sheet:

### **Emission Spectra**

Use the hand held spectroscopes to "look" at light from different sources.

What are the differences in the spectra from the various sources? Why are the spectra different?

What use can be made of the difference in spectra?

# **Exposure Levels**

#### Apparatus

tables showing typical doses of radiation, for example those below, list of recommended exposure limits

#### Action

The students examine the tables and compare the typical dosages to the recommended limits.

#### **The Physics**

Radiation comes from many natural sources, as well as man made sources, and the typical exposures are well below recommended limits. People who work with radiation such as x-ray sources, especially radiographers and dentists, and aircraft pilots who are exposed to a lot of cosmic radiation receive much higher dosages of radiation than most other people.

#### Typical average annual doses due to natural radiation.

Source	Dose (µSv / yr)
local gamma radiation	400
carbon 14	10
radon and decay products	800
potassium 40 (in body)	200
cosmic radiation	300
uranium and thorium (in body)	170

#### Typical average annual doses due to man made radiation.

Source	Dose (µSv / yr)
diagnostic radiology (eg x-rays)	220
therapeutic radiology	30
radioisotopes in medicine	2
radioactive waste	2
fallout (nuclear weapons)	10

Dosage limits for ionizing radiation (from the ARPANSA website):

The NHMRC recommended radiation dose limit for the public is 1 mSv (1000  $\mu$ Sv) per year.

ANSTO's dose constraint for reactors, which has been agreed by the Nuclear Safety Bureau, is 100  $\mu Sv$  per year for members of the public

#### Accompanying sheet:



Look at the chart showing the recommended exposure limits.

How do these compare with the dosages shown in the table?

What sort of professions do you think have the highest exposure?

## Hydrogen Spectrum

#### Apparatus

hand held spectrometer, hydrogen lamp,

#### Action

The students observe the spectral lines and use Bohr's model to explain the lines.

#### **The Physics**

The lines correspond to electron transitions, where the emitted photon has the energy difference between the initial and final energy level. This energy, and hence the wavelength, can be determined using the Bohr model for hydrogen.



A student at the Australian Catholic University observing a hydrogen spectrum.

#### Accompanying sheet:

### **Hydrogen Spectrum**

Examine the hydrogen lamp with the naked eye.

Now have a look through the spectroscope.

What do you see?

Sketch the spectra you observe.

# **Identify the Element**

#### Apparatus

hand held spectrometer, discharge lamp with label covered or removed, selection of diagrams of spectra for different elements

#### Action

The students observe the spectral lines and compare them to the diagrams given to identify the element contained in the lamp.

#### The Physics

The emission spectra for each element is unique, as they all have different energy levels. This provides a way of identifying what elements are contained in an unknown sample.





Students at the Australian Catholic University looking at the spectrum of fluorescent lights and using a chart of known spectra to identify the element (mercury) contained in the lights.

#### Accompanying sheet:

### **Identify the Element**

Use the spectroscope to look at the lamp.

Now use the diagrams of spectra from known samples to identify the element in the lamp.

What would a spectrum from a sample with several elements look like?

# Laser Light I-Focus

#### Apparatus

laser, "white light" collimated source (or a pencil torch), used blackboard dusters

#### Action

The students observe the spots from the two light sources on a distant screen (or wall). They then observe the beams from the side. The dusters can be banged together to make the laser beam visible, after students have first noticed that the laser beam is invisible from the side.

#### **The Physics**

The laser produces a small spot on the wall or screen which varies very little when the distance between the laser and screen changes. The incandescent light produces a large spot, which varies in intensity from the centre out. The spot from the incandescent light varies a lot when the distance is changed, getting larger and less intense as the distance increases. A laser beam is much more highly collimated and focused than is possible with a normal incandescent light source.





# Laser light II – Spectrum

#### Apparatus

laser, "white light" collimated source (or a torch), prism

#### Action

The students observe the resulting spectrum when light from each of the sources is passed through the prism.

#### The Physics

The incandescent light source will produce a rainbow because it acts as a blackbody and produces a continuous spectrum. The laser produces almost monochromatic light due to electron transitions.





# **Light Emitting Diodes**

#### Apparatus

a selection of LEDs, a power supply, a switch, voltmeter, variable resistor

The switch, power supply variable resistor and an LED are wired in series, with the circuit set so that the LEDs can be swapped. The voltmeter is connected across the led.

#### Action

The students look at the various LEDs in the circuit, and try to explain why they produce different colours. The students adjust the variable resistor until the LED starts to glow. They then measure voltage across the LED. They should recognize that the shorter the wavelength produced, the greater the voltage needed, and relate this to the band gap of the semiconductor material of the LED.

#### The Physics

The colour of the LED depends on the band gap. When an electron transition occurs a photon with energy depending on the change in energy of the electron is emitted.





### **Measuring Momentum and Position I**

#### Apparatus

marbles, slide, carbon paper and paper

The slide is mounted on a table such that a marble released at the top slides down and is ejected horizontally from the end of the slide. The marble falls and hits the carbon paper, which leaves a mark on the paper beneath it.

#### Action

The students release the marbles from the top of the slide, and using the mark on the paper from the carbon paper they calculate the horizontal momentum of the marble just prior to the collision. The students need to know the mass of the marble and the height of the desk.

#### **The Physics**

When the marble is released from the top of the slide it rolls down, gathering momentum as it falls. As it leaves the end of the slide it has horizontal velocity *v*. It is accelerated vertically due to gravity, and hits the floor at a time  $t = \left(\frac{2h}{g}\right)^{1/2}$  after it leaves the end of the slide, where *h* is the height of the end of the slide above floor level. The collision with the floor/paper changes the momentum of the marble, hence this measurement is of the position of the marble, and we can infer from it the momentum prior to the measurement, but not during the measurement.





### **Measuring Momentum and Position II**

#### Apparatus

marbles, slide, web cam or light gate or other passive detector

The slide is mounted on a table such that a marble released at the top slides down and is ejected horizontally from the end of the slide. The camera is used to take a picture when the marble reaches a certain height, or a light gate or some other sort of sensor which uses light to tell when the marble reaches a particular height.

#### Action

The students release the marbles from the top of the slide, and using the sensor can calculate the momentum and position of the marble at a given point.

#### **The Physics**

When the marble is released from the top of the slide it rolls down, gathering momentum as it falls. As it leaves the end of the slide it has horizontal velocity *v*. It is accelerated vertically due to gravity, and at a time  $t = \left(\frac{2h}{g}\right)^{1/2}$  after it leaves the end of the slide it has fallen a distance *h*. Knowing the horizontal distance it has traveled in this time, we can find both the marble's momentum and position. In this case the measurement has not significantly affected the momentum, however if the object were an electron, the scattering the light used to make the measurement would have had a significant effect.





### **Measuring Radiation**

#### Apparatus

TLD badges, film badges, Geiger counter or any radiation measuring devices available, sources such as smoke detectors, old gas lamp mantles, old watches with glowing numbers

#### Action

The students examine the devices and try to explain how they work. They also measure the radiation from various sources.

#### The Physics

The GM tube in the Geiger counter is filled with low pressure gas, and around +400 Volts are applied to the thin wire in the middle. When a particle enters the tube, it ionizes a gas atom. The electron is attracted to the central wire, and as it rushes towards the wire, it ionizes more gas atoms, giving an ion cascade and creating a pulse which can be amplified and counted.

Radioactivity will darken ("fog") the photographic film in a film badge. The badges have "windows" made of different materials, which block different radiation, so that the dose of  $\alpha$ ,  $\beta$  and  $\gamma$  can be distinguished.

Scintillation detectors work by the radiation striking a suitable material such as sodium iodide and causing a tiny flash of light, which is picked up by a photo-multiplier tube.





#### Accompanying sheet:

### **Measuring Radiation**

Different means of measuring radiation are shown.

Explain how they work.

Which ones would be suitable monitoring devices for persons working in a radiation area?

### **Molecular Models**

#### Apparatus

ball and stick models of atoms.

Polystyrene balls with sticks representing the outer shell electrons available for bonding. The balls could be coloured for quick identification. There should be more than one atomic species from at least a couple of periods, for example carbon and silicon or hydrogen and lithium.

#### Action

The students examine the models and try to match atoms from the same period. They should also try to explain what determines the chemical behaviour of the species.

#### **The Physics**

The chemical properties are determined by the number of outer shell electrons, which is determined by the quantum numbers, (n, l and m) and the Pauli exclusion principle. Hence atoms from the same period, which have the same number of outer-shell electrons, will have similar chemical properties.





### **Molecular Models of Semiconductors**

#### Apparatus

ball and stick models of Si, P and Al atoms.

Polystyrene balls with sticks representing the outer shell electrons available for bonding. The balls could be coloured for quick identification.

#### Action

The students examine the models and put together an *n* type and a *p* type semiconductor.

#### **The Physics**

Adding P to the Si gives an n type semiconductor, as the fifth outer shell electrons cannot be involved in bonding and hence are free to wander the lattice as charge carriers. The Al impurities give a deficit of bonding electrons, leading to holes as charge carriers.





### **Nuclear Power Stations**

#### Apparatus

diagram showing nuclear power reactor and energy transformations

#### Action

The students examine the diagram and try to understand the chain of energy transformations, and the functions of the main components.

#### The Physics

See diagram below. Note that some energy is lost as heat at various stages so most of the original energy from the nuclear process is not converted to electricity. This is true of any power plant, regardless of initial energy source.



#### Accompanying sheet:

### **Nuclear Power Stations**

Examine the diagram showing how the reactor is used to produce electricity.

In simple terms, what is the reactor used to do?

Draw a flow chart showing the conversion of energy from mass through to electricity.

# **Periodic Table**

#### Apparatus

large chart of the periodic table of the elements

#### Action

The students look at the table and try to explain the relationship between position and characteristics, such as ionization energy, bonding, etc.

#### The Physics

The elements are arranged according to their number of outer shell electrons, which is determined by the quantum numbers, (n, l and m) and the Pauli exclusion principle.



A student at the University of Sydney looking at a periodic table.

#### Accompanying sheet:

# **Periodic Table** Examine the chart of the periodic table. Why are there only 8 elements in both the second and third period? Shouldn't there be more in the third? Why do elements in a given column have similar characteristics?

# **Photoelectric Effect**

#### Apparatus

ultraviolet light source, e.g. a mercury lamp, an electroscope, a metal plate The metal plate, e.g. zinc or aluminium, is placed atop the electroscope and covered by a perspex sheet.

#### Action

The Students charge the electroscope so that it is negative, and the foil leaves move apart. The UV lamp is then turned on, and the perspex sheet removed. The electroscope leaves fall back together.

#### The physics

The UV photons collide with electrons in the metal plate, ejecting them if the energy of the photons is greater than the binding energy of the metal. The ejected electrons reduce the excess negative charge on the plate, which is connected to the electroscope. This reduces the charge on the leaves of the electroscope and they fall back together.

Note: it is almost impossible to start neutral and remove enough electrons to charge the electroscope as they recombine quickly to leave the electroscope neutral again.



Note: the perspex sheet is useful if the lamp takes a few minutes to warm up. The lamp can be turned on and the sheet used to block the UV radiation until the students are ready to use the demonstration.



### **Potential Wells and Wave Functions**

#### Apparatus

A computer simulation, such as CUPSQM, which can generate wave functions for particles in potential wells or a set of diagrams showing wave functions for particles in potential wells

#### Action

The students identify the axes, and that the wave function is superimposed over the potential well, and is **not** measured in units of energy.

#### The Physics

The potential well has a vertical axis representing potential energy, which is infinite at the edges of the well. The horizontal axis represents displacement. The wave function represents the square root of the probability density, and is **not** measured in terms of energy. It is superimposed *over the well*. Many students find these diagrams particularly abstract and difficult to interpret – particularly as there are generally no units given on the axes, and they may not be aware that this is really two plots in one.





### **Semiconductor Diode**

#### Apparatus

a germanium diode, a power supply, a switch, a small low voltage globe, a match or other heat source

The globe, switch, power supply and germanium diode are wired in series.

#### Action

The students close the switch to complete the circuit. They then heat the germanium to light the globe. They swap the polarity and repeat the process.

#### **The Physics**

When the germanium is heated more electrons move into the conduction band, allowing current to flow and the globe to light up. In reverse bias the globe takes longer to light and does not glow so brightly.



#### Accompanying sheet:

### **Semiconductor Diode**

Diodes are made from semiconductor material, such as germanium.

Use a match to heat the germanium diode, to light the lamp.

Can you light the lamp with the diode connected the other way?

Explain some uses for a diode in an electrical circuit.

### **Smoke Detector**

#### Apparatus

a battery operated smoke detector, opened up

The smoke detector should be opened up so circuitry is exposed and the warning stickers on the radioactive source visible.

#### Action

The students examine the smoke detector and identify key components including the  $^{241}$ Americium  $\alpha$  particle source.

#### The Physics

The  $\alpha$  particles ionize air molecules between two charged plates. The positive ions go to the negative plate, the negative ions to the positive plate, which gives a current. When smoke enters the space between the plates the ions attach themselves to the heavy smoke particles and the flow of current is disrupted, setting off the alarm.



#### Accompanying sheet:

### **Smoke Detector**

Examine the smoke detector. It contains a radioactive source, <sup>241</sup>Americium, which emits 5.4 MeV  $\alpha$  particles.

> The  $\alpha$  particles ionize air molecules between two charged plates. The positive ions go to the negative plate, and the negative ions to the positive plate, which gives a current.

How does smoke disrupt the current? What energy is emitted by the source per second? per day? How does this compare to acceptable exposure limits?

### **Solar Panels**

#### Apparatus

solar panel, ammeter and/or voltmeter, light source

#### Action

The students hold the cell under the light, and vary the intensity to see what effect there is on the output from the cell.

#### The Physics

The electrons absorb energy (light), allowing them to jump across the band gap, and cause a current to flow. The current is proportional to the light intensity.



#### Accompanying sheet:

### **Solar Panels**

Examine the solar panel.

How does it use photons to produce an electric current?

How is this similar to a diode?

How is it different?

### **Thermocouples and Thermistors**

#### Apparatus

a sample of semiconductor, a sample of metal, one or two ohm meters and a container of hot water to heat the samples in

A neat way to set this up is to have the samples each wired directly into the connectors of a multimeter with a large display, with the meter already set on an appropriate range.

#### Action

The students measure the resistance of each sample then heat it in the hot water and measure the resistance again.

#### The Physics

When a semiconductor is heated some of the electrons from the valence band are thermally excited into the conduction band, thus giving more charge carriers and a lower resistance. When a metal is heated it already has plenty of charge carriers in the conduction band, and the heat causes random oscillations of the lattice, leading to a shorter mean free path for the electrons and hence a greater resistance. So the sample with increasing resistance with heat is the metal, a thermocouple, the other is the thermistor.





## Transistors

#### Apparatus

transistor, power supply, voltmeter, ammeter, resistors

The components are wired up in a simple circuit, for example as an amplifier.

#### Action

The students examine the circuit and investigate the effect of varying the supply voltage using the voltmeter and ammeter.

#### **The Physics**

The transistor amplifier works by a small change in the current in the base producing a large change in current in the collector, which gives a large signal across the load resistor.



Accompanying sheet:

### Transistors

Examine the circuit. What does the transistor do?

Investigate the behaviour of the transistor by varying the voltage supplied.

What happens when you increase or decrease the voltage?

### Wave and Particle Nature of Light I: Interference Pattern

#### Apparatus

a laser, slides with single and double slits

#### Action

The students use the laser to observe interference patterns. The students are encouraged to draw diagrams showing why the interference pattern occurs, and they should explain why this shows that light has wave properties.

#### **The Physics**

Waves passing through the two slits interfere to give a pattern of fringes, with spacing depending on wavelength and slit separation. This shows the wave nature of light. Note that there are also diffraction patterns visible.



Note: This is used with both the waves workshops and the quantum workshops.



# Wave and Particle Nature of Light II: Spectral Lines

#### Apparatus

spectrometer, light source, e.g. a hydrogen lamp

#### Action

The students observe the spectral lines and try to explain them.

#### **The Physics**

Only a single discrete quantum of light, i.e. a photon, is emitted or absorbed during electron transitions. This shows the discrete or particle nature of light.



An environmental science student at the Australian Catholic University observing an emission spectrum.

#### Accompanying sheet:

### Wave and Particle Nature of Light II – Spectral Lines

Use the spectroscope to observe the spectra of the hydrogen lamp.

Why do you observe discrete lines? What does this experiment tell you about the nature of light?

### Waves on a String

#### Apparatus

taut cord or wire, oscillating driver with frequency control

The driver with amplifier is attached to one end of the cord, the other end is fixed.

#### Action

The students vary the frequency of the oscillator to produce standing wave patterns.

#### The Physics

Only certain wavelengths are possible for the standing waves;  $\lambda = 2l/n$ , where *l* is the length of the cord and n = 1, 2, 3... This gives certain discrete values of energy for each mode. This is analogous to the electron as a standing wave in the quantum model of an atom. The electron is confined to a region surrounding the nucleus, and the solutions to the Schroedinger equation are standing waves for the wave functions of the electrons, with certain discrete values of energy.





# **X-ray Diffraction**

#### Apparatus

an x-ray diffraction diagram, plot of intensity vs scattering angle, for example polonium as this is a simple cubic - a simulated diffraction pattern for polonium is given below, a model of the crystal - more than a single unit cell helps for multiple plane spacings to be clear.

NaCl is also good because of its everyday familiarity, however it is a face centred cubic hence has a more complex structure and diffraction pattern.

#### Action

The students examine the diffraction data and try to match the peaks to planes in the crystal, and find lattice spacings.

#### **The Physics**

The atoms in the crystal form planes, with spaces between them like the lines of a diffraction grating. The larger spacings give smaller angles, because for constructive interference  $d\sin\theta = n\lambda$ , so the peaks at larger angles are for the smaller lattice spacings, such as the planes which make the sides of the unit cells. The smaller angles are for large lattice spacings.

This is reflected beam technique, in contrast to the way x-rays are used to make x-ray pictures of bones, in which the attenuated transmitted beam is used to get information about the material.



#### Accompanying sheet:

### **X-ray Diffraction**

Examine the x-ray diffraction pattern.

Can you match the peaks to particular planes of atoms in the lattice?

Why are there peaks at multiples of the scattering angles?

# **X-ray Pictures**

#### Apparatus

two or three x-ray pictures

#### Action

The students examine the pictures and identify areas of high and low x-ray absorption, and high and low tissue density.

#### The Physics

X-rays are more strongly absorbed by denser tissues, such as bone. This is because these tissues have a higher proportion of elements with a large number of electrons, which the x-rays interact strongly with. These areas show up white on typical x-ray images, which are negatives.



#### Accompanying sheet:

### **X-ray Pictures**

Examine the x-ray films.

Why are some areas light and others dark? In which areas are more x-rays absorbed?

Why are more x-rays absorbed in these regions?

# X-ray Tube

#### Apparatus

Exposed x-ray tube, or opened up x-ray machine showing target and electron source.

#### Action

The students examine the tube and draw a diagram showing the main components.

#### **The Physics**

Electrons are emitted from the filament, and accelerated towards the target. At the target they collide with atoms of the target material and x-ray photons are emitted.



Students at the University of Sydney discussing the workings of an x-ray tube.

#### Accompanying sheet:

### X-ray Tube

Examine the x-ray tube and draw a diagram showing the main components.

Where is the target?

Where and how are the electrons accelerated?