

Lecture 10

# Work, Power and Potential energy

## Energy

kinetic energy – energy of motion

$$K = \frac{1}{2} mv^2$$

potential energy – stored energy

$$U = mgh \quad (\text{gravity})$$

Kinetic energy and potential energy added together are called *Mechanical Energy*.

KJF §10.5

2

## Law of Conservation of Energy

Energy cannot be created or destroyed  
(i.e. it is "conserved")

It can only be changed from one form to another

OR

In an isolated system — one where there is no  
energy transfer into or out of the system — the  
total energy  $E_{\text{tot}}$  is conserved.

KJF §10.3

3


## WORK

KJF §10.4

4

## What is Work?

Work is the process of transferring energy from  
the environment to a system, or from the system to  
the environment, by the application of forces.



KJF §10.2

5

## What is Work?

“Doing work” means using a **force** to

- transfer energy from one object to another, or
- convert energy from one form to another.

Work ( $W$ ) is equal to the **amount of energy**  
transferred or converted by the force.

Work is a scalar. S.I. unit is also the joule (J).

If force is constant then


$$W = Fs \cos \theta$$

where  $F$  is applied force,  $s$  is object's displacement while  
the force is applied and  $\theta$  is angle between applied force  
and displacement.

6


## Work and Kinetic Energy (1)

Consider a single constant force  $F$  acting on a body causing a change in kinetic energy only. Suppose it moves in the  $x$ -direction and the force acts at angle  $\theta$  to that direction.



KJF §10.5

7



If body moves displacement  $d = x_f - x_i$ . From the motion equations;

$$v_f^2 = v_i^2 + 2a_x d \quad \text{where } d = x_f - x_i$$

$$\text{i.e. } 2a_x d = (v_f^2 - v_i^2)$$

$$\begin{aligned} \text{But the K.E. is changing; } \Delta K &= K_f - K_i \\ &= \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2 = \frac{1}{2}m(v_f^2 - v_i^2) = \frac{1}{2}m \times 2a_x d \end{aligned}$$

KJF §10.5

8

## Work and Kinetic Energy (2)

$$\begin{aligned} \text{Therefore } \Delta K &= \frac{1}{2}m \times 2a_x d = ma_x d = F_x d \\ &= F \cos \theta d \end{aligned}$$

But our definition of work was “change in energy” and in this case the only change is in kinetic energy.

Therefore the work done is

$$W = \Delta K = Fd \cos \theta$$

KJF §10.4

9

## Sign Convention for Work

- **Work can be negative!** (i.e. Energy will be transferred *from* the body). (“signed scalar”)
- If the applied force is causing the object to **increase** in energy, then the work is **positive**, e.g. work done by legs walking upstairs.
- If the applied force is causing the object to **decrease** in energy, then work is **negative**, e.g. work done by legs walking downstairs.

10

## Forces that do no work

- If the object does not move, no work is done




Copyright © 2007, Pearson Education, Inc., publishing as Pearson Addison-Wesley

KJF §10.4

11

## Forces that do no work

- If the force is perpendicular to the displacement, then the force does no work



Copyright © 2007, Pearson Education, Inc., publishing as Pearson Addison-Wesley

KJF §10.4

12

## Work–Energy Theorem

Work is a scalar, i.e. the work done by the individual forces can be added together "arithmetically":

$$W = W_1 + W_2 + \dots$$

**Work-Energy Theorem:** (a special case of the law of conservation of energy) The change in kinetic energy of a system equals the sum of work done by all the individual forces on the system:

$$\Delta K = \sum W$$

*Strictly speaking this theorem applies to rigid ("non-squishy") objects*

KJF §10.3

13

## Example: KJF problem 10.11

A 20 g plastic ball is moving to the left at 30 m/s. How much work must be done on the ball to cause it to move to the right at 30 m/s?



[0 J]

14

## What is Power?

**Power** means the **rate** at which a force does work on an object.

Instantaneous power:  $P = \frac{dW}{dt}$

Average  $P = \frac{\text{work done}}{\text{time taken}} = \frac{\Delta W}{\Delta t}$

Also remember that  $W = Fs \cos\theta$ , so if  $F$  and  $\theta$  are constant

$$P = \frac{\Delta W}{\Delta t} = F \frac{\Delta s}{\Delta t} \cos\theta = Fv \cos\theta = \mathbf{F} \cdot \mathbf{v}$$

KJF §10.10

15

## Units

Units: watt (W) = joule.second<sup>-1</sup> (J.s<sup>-1</sup>)

[1 horsepower (hp) = 746 W]

**Note!** Do not confuse the unit watt (W) with the algebraic symbol for work (W).

*The 1st is the unit for power P, the 2nd represents an energy and has joules (J) as its unit.*

	Algebraic Symbol	Unit Symbol
Energy, Work	K, U, W...	J
Power	P	W

16

## Problem

The loaded cab of an elevator has a mass of  $3.0 \times 10^3$  kg and moves 210m up the shaft in 23s at constant speed.

At what average rate does the force from the cable do work on the cab?

[ $2.7 \times 10^5$  W]

17

## Gravitational Potential



Two bodies are attracted because of gravity caused by their mass, so we have to do work  $W$  to push them apart.

When we do this, energy is transferred to the system — called **gravitational potential energy**,  $U$ . Near earth's surface  $U = mgh$

$U$  is a property of the **whole system** e.g. system is earth + ball, but usually we simplify and say "the ball has potential energy"

KJF §10.6

18

Work done **by** weight (gravity) on the ball as it falls a distance  $h$  towards earth is

$$W = Fs \cos \theta = mg \times \Delta h \times \cos 0^\circ = mg\Delta h$$

*i.e.* work done by weight (gravity) = minus the change in  $U$



Only the **change** in height is significant  
 $\Rightarrow$  we are free to choose any reference level we like where  $U=0$




19

## PE is independent of path

Because gravitational potential energy depends only on the height of the object above the reference level, the potential energy does **not** depend on the path taken.

In both cases the G.P.E increased by the same amount:  $mgh$




KJF §10.6

20



## Conservative Forces

If work done by a force in moving an object from A to B does NOT depend on the path taken, we call it a **conservative force** (e.g. gravity, ideal springs).



An object moving under the influence of a conservative force always conserves mechanical energy:

$$ME = K_i + U_i = K_f + U_f$$


EVERY conservative force has "potential energy" associated with it (e.g. gravity  $\rightarrow$  gravitational P.E., spring force  $\rightarrow$  elastic P.E.).

KJF §10.6

21

Forces like friction which dissipate energy instead of storing it are **non-conservative forces**.

If an object moves in a closed loop under the influence of a conservative force, the total work done is **ZERO**.



Work done in moving an object from A  $\rightarrow$  B under the influence of a conservative force is the exact negative of work done in going from B  $\rightarrow$  A, i.e. **reversible**

$$W_{AB} = -W_{BA}$$

e.g. Work done by gravity going upstairs and back downstairs again.

22